

For Research Use Only

ARNT/HIF-1 beta Monoclonal antibody

Catalog Number: 66732-1-Ig



Basic Information

Catalog Number:

66732-1-Ig

Size:

150ul, Concentration: 1300 ug/ml by 405

Nanodrop and 1000 ug/ml by Bradford

method using BSA as the standard;

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG5507

GenBank Accession Number:

BC060838

GeneID (NCBI):

405

UNIPROT ID:

P27540

Full Name:

aryl hydrocarbon receptor nuclear translocator

Calculated MW:

416 aa, 46 kDa, 87 kDa

Observed MW:

87 kDa

Purification Method:

Protein G purification

CloneNo.:

3B12C5

Recommended Dilutions:

WB: 1:2000-1:10000

IF/ICC: 1:650-1:2600

FC (Intra): 0.50 ug per 10⁶ cells in a 100 µl suspension

Applications

Tested Applications:

WB, IF/ICC, FC (Intra), ELISA

Species Specificity:

human, mouse, rat

Positive Controls:

WB: Caco-2 cells, HeLa cells, HepG2 cells, HT-29 cells, HSC-T6 cells, NIH/3T3 cells

IF/ICC: HT-1376 cells,

FC (Intra): HeLa cells,

Background Information

HIF1B(HIF-1 beta), also named as ARNT, is a 789 amino acid protein, which forms a heterodimer with AHR, AHRR, HIF1A and EPAS1/HIF2A as well as with other bHLH proteins. HIF1B is Required for activity of the Ah (dioxin) receptor. This protein is required for the ligand-binding subunit to translocate from the cytosol to the nucleus after ligand binding. The heterodimer with HIF1A or EPAS1/HIF2A functions as a transcriptional regulator of the adaptive response to hypoxia

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 66732-1-Ig (ARNT/HIF-1 beta antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HT-1376 cells using ARNT/HIF-1 beta antibody (66732-1-Ig, Clone: 3B12C5) at dilution of 1:1300 and Multi-rAb CoraLite® Plus 594-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (Cat.NO. RGAM004).



1X10⁶ HeLa cells were intracellularly stained with 0.5 ug Anti-Human ARNT/HIF-1 beta (66732-1-Ig, Clone:3B12C5) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.5 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).