

For Research Use Only

CHOP; GADD153 Monoclonal antibody, PBS Only

Catalog Number: 66741-1-PBS



Basic Information

Catalog Number:

66741-1-PBS

Size:

100ug, Concentration: 1 mg/ml by
Nanodrop;

Source:

Mouse

Isotype:

IgG2a

Immunogen Catalog Number:

AG7354

GenBank Accession Number:

BC003637

GeneID (NCBI):

1649

UNIPROT ID:

P35638

Full Name:

DNA-damage-inducible transcript 3

Calculated MW:

19 kDa

Observed MW:

30 kDa

Purification Method:

Protein A purification

CloneNo.:

4F3G1

Applications

Tested Applications:

WB, IHC, Indirect ELISA

Species Specificity:

Human, mouse, rat

Background Information

CHOP, also known as GADD153 or DDIT3, is a highly conserved gene in both the structural and regulatory regions. Induced by unfolded and misfolded proteins, CHOP is significantly induced by ER stress. CHOP is considered a proapoptotic marker of ER stress dependent cell death. CHOP acts as a dominant-negative inhibitor of the transcription factor C/EBP and LAP. It may play an important role in the malignant transformation of nevus to melanoma. The calculated molecular weight of CHOP is 19 kDa, but the protein migrates on an SDS-PAGE gel with an observed molecular mass of 29 kDa (PMID: 1547942).

Storage

Storage:

Store at -80°C.

Storage Buffer:

PBS only


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.



Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 66741-1-Ig (CHOP; GADD153 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66741-1-PBS in a different storage buffer formulation.



Un-treated and Tunicamycin treated HepG2 lysates were subjected to SDS PAGE followed by western blot with 66741-1-Ig (CHOP; GADD153 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66741-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human cervical cancer tissue slide using 66741-1-Ig (CHOP; GADD153 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66741-1-PBS in a different storage buffer formulation.