

## Factor X Monoclonal antibody

Catalog Number: 66753-1-Ig

## Basic Information

<b>Catalog Number:</b> 66753-1-Ig	<b>GenBank Accession Number:</b> BC046125	<b>Purification Method:</b> Protein G purification
<b>Size:</b> 150ul , Concentration: 1800 ug/ml by Nanodrop and 1000 ug/ml by Bradford method using BSA as the standard;	<b>GeneID (NCBI):</b> 2159	<b>CloneNo.:</b> 3D3F11
<b>Source:</b> Mouse	<b>UNIPROT ID:</b> P00742	<b>Recommended Dilutions:</b> WB 1:5000-1:50000 IHC 1:500-1:2000 IF-P 1:200-1:800
<b>Isotype:</b> IgG1	<b>Full Name:</b> coagulation factor X	
<b>Immunogen Catalog Number:</b> AG11664	<b>Calculated MW:</b> 488 aa, 55 kDa	
	<b>Observed MW:</b> 50 kDa	

## Applications

**Tested Applications:**  
WB, IHC, IF-P, ELISA

**Species Specificity:**  
human

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

**Positive Controls:**

**WB :** human plasma tissue,  
**IHC :** human liver tissue,  
**IF-P :** human liver tissue,

## Background Information

Coagulation factor X (F10) is a vitamin K-dependent glycoprotein that participates in the middle phase of the blood coagulation cascade (PMID: 1931959; 1313796). It is synthesized as a single-chain precursor in the liver where it undergoes a number of covalent processing steps before secretion into the blood as a two-chain molecule linked by a disulfide bond (PMID: 1313796). The light chain contains 2 EGF-like domains, while the heavy chain contains the catalytic domain which is structurally homologous to those of the other hemostatic serine proteases. Factor X is activated into factor Xa, by both the extrinsic and intrinsic pathway. The activated factor then converts prothrombin to thrombin in the presence of factor Va, calcium and phospholipid during blood clotting.

## Storage


**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol, pH7.3  
 Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:  
 T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
 E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
 W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




human plasma were subjected to SDS PAGE followed by western blot with 66753-1-Ig (Factor X antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 66753-1-Ig (Factor X antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 66753-1-Ig (Factor X antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human liver tissue using Factor X antibody (66753-1-Ig, Clone: 3D3F11) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).