

For Research Use Only

# AGR2 Monoclonal antibody, PBS Only

Catalog Number: 66768-1-PBS



## Basic Information

<b>Catalog Number:</b> 66768-1-PBS	<b>GenBank Accession Number:</b> BC015503	<b>Purification Method:</b> Protein A purification
<b>Size:</b> 100ug , Concentration: 1mg/ml by Nanodrop;	<b>GeneID (NCBI):</b> 10551	<b>CloneNo.:</b> 1A8A8
<b>Source:</b> Mouse	<b>UNIPROT ID:</b> O95994	
<b>Isotype:</b> IgG2b	<b>Full Name:</b> anterior gradient homolog 2 (Xenopus laevis)	
<b>Immunogen Catalog Number:</b> AG2919	<b>Calculated MW:</b> 175 aa, 20 kDa	
	<b>Observed MW:</b> 17 kDa	

## Applications

**Tested Applications:**  
WB, IHC, IF/ICC, IF-P, FC (Intra), Indirect ELISA

**Species Specificity:**  
human, pig

## Background Information

AGR2, also named AG2 or HPC8, encodes anterior gradient protein 2 homolog which belongs to the AGR family. It is a secreted protein localized in endoplasmic reticulum. AGR2 plays roles in MUC2 post-transcriptional synthesis, secretion and production of mucus by intestinal cells. AGR2 was significantly elevated in the pancreatic juice from patients with pre-malignant conditions as well as pancreatic cancer compared to control pancreatic juice samples. AGR2 levels in pancreatic juice could potentially be used to aide in assessment of high-risk patients undergoing endoscopic procedures.

## Storage

**Storage:**  
Store at -80°C.

**Storage Buffer:**  
PBS Only


For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




pig stomach tissue were subjected to SDS PAGE followed by western blot with 66768-1-Ig (AGR2 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66768-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66768-1-Ig (AGR2 antibody) at dilution of 1:300 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66768-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66768-1-Ig (AGR2 antibody) at dilution of 1:300 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66768-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed A549 cells using AGR2 antibody (66768-1-Ig, Clone: 1A8A8 ) at dilution of 1:1500 and Multi-rAb CoraLite® Plus 594-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (Cat.NO. RGAM004). This data was developed using the same antibody clone with 66768-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed HT-29 cells using AGR2 antibody (66768-1-Ig, Clone: 1A8A8 ) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66768-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed human colon cancer tissue using AGR2 antibody (66768-1-Ig, Clone: 1A8A8 ) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66768-1-PBS in a different storage buffer formulation.



1X10<sup>6</sup> HT-29 cells were intracellularly stained with 0.2 ug Anti-Human AGR2 (66768-1-Ig, Clone:1A8A8) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), and 0.2 ug Mouse IgG2b Isotype Control (66360-3-Ig, Clone: K11B8C4B5) (blue). Cells were fixed with 4% PFA and permeabilized with 0.1% TritonX-100. This data was developed using the same antibody clone with 66768-1-PBS in a different storage buffer formulation.