For Research Use Only

TOM20 Monoclonal antibody, PBS Only



Catalog Number: 66777-1-PBS

Basic Information

Catalog Number:

66777-1-PBS

100ug, Concentration: 1mg/ml by

Nanodrop: Mouse

Isotype: lgG2b

Immunogen Catalog Number:

AG2378

ELISA, IF/ICC, IHC, WB

Tested Applications:

Species Specificity:

Background Information

TOM20, also named as KIAA0016, belongs to the Tom20 family. It is a central component of the receptor complex responsible for the recognition and translocation of cytosolically synthesized mitochondrial preproteins. Together $with \, TOM22, \, TOM20 \, functions \, as \, the \, transit \, peptide \, receptor \, at \, the \, surface \, of \, the \, mitochondrion \, outer \, membrane \, the \, surface \, of \, the \, mitochondrion \, outer \, membrane \, the \, surface \, of \, the \, mitochondrion \, outer \, membrane \, the \, surface \, of \, the \, mitochondrion \, outer \, membrane \, the \, surface \, of \, the \, mitochondrion \, outer \, membrane \, the \, surface \, of \, the \, mitochondrion \, outer \, membrane \, the \, surface \, of \, the \, mitochondrion \, outer \, membrane \, the \, surface \, of \, the \, mitochondrion \, outer \, membrane \, the \, surface \, of \, the \, mitochondrion \, outer \, membrane \, the \, surface \, of \, the \,$ and facilitates the movement of preproteins into the TOM40 translocation pore. TOM20 is characterized as major docking receptors to mediate the recognition by different mechanisms.

GenBank Accession Number:

translocase of outer mitochondrial

membrane 20 homolog (yeast)

BC000882

GeneID (NCBI):

UNIPROT ID: Q15388

Full Name:

Calculated MW:

145 aa, 16 kDa Observed MW: 16 kDa

Storage


Applications

Storage: Store at -80°C. Storage Buffer: PBS Only


Purification Method: Protein A purification

CloneNo.: 1D6F5


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66777-1-1g (TOM20 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66777-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 66777-1-lg (TOM20 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66777-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using TOM20 antibody (66777-1-lg, Clone: 1D6F5) at dilution of 1:800 and CoraLite® 488-Conjugated Affini Pure Goat Anti-Mouse IgG(H+L), CL594-phalloidin (red). This data was developed using the same antibody clone with 66777-1-PBS in a different storage buffer formulation.



1X10^6 HeLa cells were intracellularly stained with 0.4 ug Anti-Human TOM20 (66777-1-1g, Clone:1D6F5) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgC(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 66777-1-PBS in a different storage buffer formulation.