For Research Use Only

DIO2 Monoclonal antibody, PBS Only

Catalog Number: 66813-1-PBS

Featured Product



Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method: Protein A purification

66813-1-PBS

BC074882 GeneID (NCBI):

Size:

CloneNo.: 1F2A1

100ug, Concentration: 1 mg/ml by Nanodrop;

UNIPROT ID:

Mouse

Q92813 Full Name:

Isotype:

deiodinase, iodothyronine, type II

IgG2a

Calculated MW: 273 aa, 31 kDa

Immunogen Catalog Number: AG24057

Observed MW:

30-35 kDa

Applications

Tested Applications:

WB, IHC, ELISA

Species Specificity:

Human

Background Information

Type II iodothyronine deiodinase(DIO2), belongs to the iodothyronine deiodinase family, with two isoform of 30 and 34 kDa. DIO2 can activates thyroid hormone by converting the prohormone thyroxine (T4) by outer ring deiodination (ORD) to bioactive 3,3',5-triiodothyronine (T3). DIO2 is thought to be responsible for the 'local' production of T3, and thus important in influencing thyroid hormone action in these tissues.

Storage


Storage:

Store at -80°C.


Storage Buffer:

PBS only


Selected Validation Data




WB result of DIO2 antibody (66813-1-lg; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-DIO2 transfected MCF-7 cells. This data was developed using the same antibody clone with 66813-1-PBS in a different storage buffer formulation.



T-47D cells were subjected to SDS PAGE followed by western blot with 66813-1-1g (DIO2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66813-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human thyroid cancer tissue slide using 66813-1-lg (DIO2 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66813-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human thyroid cancer tissue slide using 66813-1-1g (DIO 2 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66813-1-PBS in a different storage buffer formulation.