

For Research Use Only

PRDX1 Monoclonal antibody

Catalog Number: 66820-1-Ig

Featured Product

5 Publications



Basic Information

Catalog Number: 66820-1-Ig	GenBank Accession Number: BC007063	Purification Method: Protein A purification
Size: 150ul, Concentration: 1000 ug/ml by Nanodrop;	GeneID (NCBI): 5052	CloneNo.: 2B2A2
Source: Mouse	UNIPROT ID: Q06830	Recommended Dilutions: WB 1:5000-1:50000 IHC 1:1000-1:4000 IF-P 1:200-1:800
Isotype: IgG1	Full Name: peroxiredoxin 1	
Immunogen Catalog Number: AG8821	Calculated MW: 199 aa, 22 kDa	
	Observed MW: 23 kDa	

Applications

Tested Applications:

WB, IHC, IF-P, ELISA

Cited Applications:

WB, IF, IP, ColP

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HeLa cells, U2OS cells, HEK-293 cells, Caco-2 cells, MCF-7 cells, Jurkat cells, HSC-T6 cells, NIH/3T3 cells, 4T1 cells, LNCaP cells

IHC: human liver cancer tissue,

IF-P: human liver cancer tissue,

Background Information

PRDX1 (Peroxiredoxin-1), also named as PAGA, PAGB, TDPX2, PAG or NKEF-A, belongs to the ahpC/TSA family. PRDX1 is a thiol reductase that plays critical roles in oxidative and thermal stress defense mechanisms through its abilities to metabolize H₂O₂ and act as a molecular chaperone, respectively. PRDX1 might participate in the signaling cascades of growth factors and tumor necrosis factor- α by regulating the intracellular concentrations of H₂O₂ (PMID: 9497357). It reduces an intramolecular disulfide bond in GDPD5 that gates the ability to GDPD5 to drive postmitotic motor neuron differentiation. PRDX1 can form a dimer, and also can be phosphorylated on Thr-90 during the M-phase, which leads to a more than 80% decrease in enzymatic activity (PMID: 22583657, 11986303).

Notable Publications

Author	Pubmed ID	Journal	Application
Ling Zhu	36290756	Antioxidants (Basel)	WB
Lin Lv	34260286	J Virol	WB, ColP
Xiang-Yu Ma	38780721	Mol Neurobiol	IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66820-1-Ig (PRDX1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




U2OS (WT and PRDX1 KO) lysates prepared with RIPA buffer, 10 µg protein loaded. 66820-1-Ig incubated at 1:5000 at 4°C overnight in 5% BSA in TBST. Ponceau stained transfers shown on right. Data provided by YCharOS, an open science company with a mission to validate commercial antibodies to improve scientific reproducibility and transparency.




WB result of PRDX1 antibody (66820-1-Ig; 1:200000; incubated at room temperature for 1.5 hours) with sh-Control and sh-PRDX1 transfected HEK-293 cells.




Immunofluorescent analysis of (4% PFA) fixed human liver cancer tissue using PRDX1 antibody (66820-1-Ig, Clone: 2B2A2) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Various lysates were subjected to SDS PAGE followed by western blot with 66820-1-Ig (PRDX1 antibody) at dilution of 1:50000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 66820-1-Ig (PRDX1 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 66820-1-Ig (PRDX1 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).