For Research Use Only

## human IBA1 Monoclonal antibody

Catalog Number:66827-1-lg 47 Publications



Basic Information	Catalog Number: 66827-1-lg	GenBank Accession Number: BC009474	Purification Method: Protein A purification	
	Size:	GeneID (NCBI):	CloneNo.:	
	150ul, Concentration: 2144 ug/ml by		1C6A10	
	Nanodrop and 1000 ug/ml by Bradford method using BSA as the standard;	ENSEMBL Gene ID: ENSG00000204472	Recommended Dilutions: IHC 1:1000-1:4000	
	Source:	UNIPROT ID:	IF-Fro 1:200-1:800	
	Mouse	P55008		
	Isotype: IgG2a	Full Name: allograft inflammatory factor 1		
	Immunogen Catalog Number: AG28236	Calculated MW: 17 kDa		
Applications	Tested Applications: Positive Controls:		ive Controls:	
	IHC, IF-Fro, ELISA Cited Applications:		: human brain tissue, human gliomas tissue, Ian liver cancer tissue, human tonsillitis tissue	
	WB, IHC, IF IF-Fro : mouse brain tissue, Species Specificity: human		o : mouse brain tissue,	
	Cited Species: human			
	Note-IHC: suggested antigen r			
	TE buffer pH 9.0; (*) Alternativ retrieval may be performed w buffer pH 6.0			
Background Information	retrieval may be performed w buffer pH 6.0 IBA1 is a 143 amino acid cytoplasmic monocytes and macrophages and is k	<b>ith citrate</b> ;, inflammation response scaff nown to be involved in macro -regulated in activated micro	fold protein. It is constitutively expressed in phage activation. It is a marker of activated glia following facial nerve axotomy, ischemi phenotypes of microglia.	
	retrieval may be performed w buffer pH 6.0 IBA1 is a 143 amino acid cytoplasmic monocytes and macrophages and is k macrophage. Expression of IBA1 is up and several brain diseases, thereby in	<b>ith citrate</b> ;, inflammation response scaff nown to be involved in macro -regulated in activated micro	pphage activation. It is a marker of activated glia following facial nerve axotomy, ischemi	
	retrieval may be performed w buffer pH 6.0 IBA1 is a 143 amino acid cytoplasmic monocytes and macrophages and is k macrophage. Expression of IBA1 is up and several brain diseases, thereby in Author Pubr	ith citrate , inflammation response scaft nown to be involved in macro -regulated in activated micro nplicating it in the activated p	ophage activation. It is a marker of activated glia following facial nerve axotomy, ischemi phenotypes of microglia. Application	
	retrieval may be performed w buffer pH 6.0 IBA1 is a 143 amino acid cytoplasmic monocytes and macrophages and is k macrophage. Expression of IBA1 is up and several brain diseases, thereby in Author Pubr Yingchu Gu 3447	ith citrate , inflammation response scaft nown to be involved in macro -regulated in activated micro nplicating it in the activated p ned ID Journal	pphage activation. It is a marker of activated glia following facial nerve axotomy, ischemi phenotypes of microglia. Application i IF	
Background Information Notable Publications	retrieval may be performed w buffer pH 6.0 IBA1 is a 143 amino acid cytoplasmic monocytes and macrophages and is k macrophage. Expression of IBA1 is up and several brain diseases, thereby in Author Pubr Yingchu Gu 3447 Shiji Huo 3483	ith citrate ;, inflammation response scaff nown to be involved in macro -regulated in activated micro nplicating it in the activated p ned ID Journal /1984 J Mol Neurosci	phage activation. It is a marker of activated glia following facial nerve axotomy, ischemi phenotypes of microglia. Application i IF mation IF	

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Immunohistochemical analysis of paraffinembedded human brain tissue slide using 66827-1-Ig (IBA1 antibody) at dilution of 1:4000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human brain tissue slide using 66827-1-Ig (IBA1 antibody) at dilution of 1:4000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66827-1-1g (IBA1 antibody) at dilution of 1:1000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66827-1-lg (IBA1 antibody) at dilution of 1:1000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed frozen OCT-embedded mouse brain tissue using human IBA1 antibody (66827-1-Ig, Clone: 1C6A10) at dilution of 1:400 and Coralite®488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1).