For Research Use Only

CaMKII alpha Monoclonal antibody

Catalog Number: 66843-1-lg 4 Publications



Basic Information

Catalog Number: GenBank Accession Number:

66843-1-Ig BC040457
Size: Genel D (NCBI):
150ul , Concentration: 1300 ug/ml by 815

Nanodrop and 1000 ug/ml by Bradford UNIPROT ID: method using BSA as the standard; Q9UQM7

Source: Full Name:

Mouse calcium/calmodulin-dependent

Isotype: protein kinase II alpha IgG1 Calculated MW:

Immunogen Catalog Number: 478 aa, 54 kDa AG4736 Observed MW: 45-60 kDa

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA Cited Applications:

WB, IF

Species Specificity: Human, Mouse, Rat, Pig

Cited Species: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: pig brain tissue, pig cerebellum tissue, rat brain tissue, rat cerebellum tissue, mouse brain tissue, mouse cerebellum tissue, rabbit brain tissue, rat rain

Purification Method:

Protein A purification

Recommended Dilutions:

WB 1:5000-1:50000

IF/ICC 1:200-1:800

IHC 1:500-1:2000

CloneNo.:

1E3B3

IHC: mouse brain tissue, human renal cell carcinoma

tissue

IF/ICC: SH-SY5Y cells,

Background Information

CaM-kinase II (CAMK2) is a prominent kinase in the central nervous system that may function in long-term potentiation and neurotransmitter release. It is a member of the NMDAR signaling complex in excitatory synapses it may regulate NMDAR-dependent potentiation of the AMPAR and synaptic plasticity. This antibody can recognize all the members of the CAMK2 family.

Notable Publications

| Author | Pubmed ID | Journal | Application |
|----------------|-----------|-------------------|-------------|
| Ting-Ting Chen | 33714957 | Aging (Albany NY) | WB |
| Weili Wang | 39896347 | Noncoding RNA Res | WB |
| Yan Sun | 38852473 | Phytomedicine | WB |

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




Immunofluorescent analysis of (4% PFA) fixed SH-SY5Y cells using CAMK2 antibody (66843-1-Ig, Clone: 1E3B3) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Mouse IgG(H+L).




Various lysates were subjected to SDS PAGE followed by western blot with 66843-1-lg (CAMK2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 66843-1-Ig (CAMK2 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 66843-1-Ig (CAMK2 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 66843-1-lg (CAMK2A antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.