For Research Use Only

SPON2 Monoclonal antibody

Catalog Number:66859-1-lg 1 Publications


Antibodies | ELISA kits | Proteins www.ptglab.com


Basic Information	Catalog Number: 66859-1-lg	GenBank Accession Number: BC002707	Purification Method: Protein A purification
	Size: 150ul, Concentration: 1500 ug/ml by Nanodrop and 1000 ug/ml by Bradforc method using BSA as the standard; Source: Mouse Isotype: IgG2a Immunogen Catalog Number: AG14337	GenelD (NCBI): 10417	CloneNo.: 1H4B11 Recommended Dilutions: WB 1:1000-1:4000 IHC 1:500-1:2000
		Full Name: spondin 2, extracellular matrix protein Calculated MW: 331 aa, 36 kDa Observed MW: 38 kDa	
Applications	Tested Applications: WB, IHC, ELISA	Positive Controls: WB : LNCaP cells, IHC : human prostate cancer tissue,	
	Cited Applications: WB, IHC		
	Species Specificity: Human		
	Cited Species: human		
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0		
Background Information	SPON2 (Spondin 2), also known as Mindin, is a member of the F-spondin family of secreted extracellular matrix proteins. This protein has been implicated in axon guidance, immune response and adhesion. SPON2 is essential in the initiation of the innate immune response and represents a unique pattern-recognition molecule in the ECM for microbial pathogens (PMID: 14691481). It also functions as an integrin ligand for inflammatory cell recruitment and T-cell priming (PMID: 19153605). Besides, SPON2 has been reported as a prognostic biomarker for ovarian and prostate cancer (PMID: 17490732; 22615945; 23665271).		
5	microbial pathogens (PMID: 14691481 T-cell priming (PMID: 19153605). Besi	des, SPON2 has been reported as	ligand for inflammatory cell recruitment and
Notable Publications	microbial pathogens (PMID: 14691481 T-cell priming (PMID: 19153605). Besi prostate cancer (PMID: 17490732; 226	des, SPON2 has been reported as	ligand for inflammatory cell recruitment and
	microbial pathogens (PMID: 14691481 T-cell priming (PMID: 19153605). Besi prostate cancer (PMID: 17490732; 226 Author Pubr	des, SPON2 has been reported as 15945; 23665271).	ligand for inflammatory cell recruitment an a prognostic biomarker for ovarian and
	microbial pathogens (PMID: 14691481 T-cell priming (PMID: 19153605). Besi prostate cancer (PMID: 17490732; 226 Author Pubr	des, SPON2 has been reported as 15945; 23665271). med ID Journal 27776 Bone er shipment. % glycerol pH 7.3.	ligand for inflammatory cell recruitment an a prognostic biomarker for ovarian and Application

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





LNCaP cells were subjected to SDS PAGE followed by western blot with 66859-1-1g (SPON2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 66859-1-1g (SPON2 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 66859-1-1g (SPON2 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).