### For Research Use Only

# Glucocorticoid receptor Monoclonal antibody



Catalog Number:66904-1-lg 5 Publications

**Basic Information** 

Catalog Number: GenBank Accession Number: Purification Method: 66904-1-lg BC015610 Protein G purification

Size:GeneID (NCBI):CloneNo.:150ul , Concentration: 1000 ug/ml by 29081D9A9

Bradford method using BSA as the standard; P04150 Recommended Dilutions:

P04150 WB 1:20000-1:100000

Source: Full Name: IHC 1:150-1:600

Mouse nuclear receptor subfamily 3, group C, IF-P 1:200-1:800

Isotype: member 1 (glucocorticoid receptor)

IgG1 Calculated MW:
Immunogen Catalog Number: 86 kDa

AG28431 Observed MW:

97 kDa

**Applications** 

Tested Applications:
WB, IHC, IF-P, ELISA
Cited Applications:
WB, IHC, IF

Species Specificity: Human, mouse, rat Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

#### **Positive Controls:**

WB: HEK-293 cells, HepG2 cells, HeLa cells, HSC-T6 cells, NIH/3T3 cells, 4T1 cells, MCF-7 cells, Jurkat cells, K-562 cells, PC-12 cells

IHC: mouse brain tissue,
IF-P: mouse brain tissue,

# **Background Information**

Glucocorticoid receptor (GR, or GCR) also known as NR3C1 (nuclear receptor subfamily 3, group C, member 1) is a receptor for glucocorticoids, which owns a dual mode of action: as a transcription factor that binds to glucocorticoid response elements (GRE) and as a modulator of other transcription factors. It is involved in cell proliferation and differentiation and specifically implicated in newborn birth weight, thus providing a biological mechanism by which NR3C1 expression may influence birth weight (PMID:22810058).

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Qi Liu	35883709	Antioxidants (Basel)	IF
Lisa M Cooper	32639872	Am J Physiol Cell Physiol	WB
Haihua Wang	39820491	Cell Death Dis	WB,IHC

Storage

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


\*\*\* 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 66904-1-1g (Glucocorticoid receptor antibody) at dilution of 1:50000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 66904-1-Ig (Glucocorticoid receptor antibody) at dilution of 1:300 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using Glucocorticoid receptor antibody (66904-1-lg, Clone: 1D9A9) at dilution of 1:400 and Coralite®488-Conjugated Goat Anti-Mouse IgG(H+L).