

For Research Use Only

DDX21 Monoclonal antibody

Catalog Number: 66925-1-Ig



Basic Information

Catalog Number: 66925-1-Ig	GenBank Accession Number: BC008071	Purification Method: Protein A purification
Size: 150ul, Concentration: 1300 µg/ml by Nanodrop and 773 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 9188	CloneNo.: 1B9C8
Source: Mouse	Full Name: DEAD (Asp-Glu-Ala-Asp) box polypeptide 21	Recommended Dilutions: WB 1:20000-1:100000 IHC 1:250-1:1000 IF 1:50-1:500
Isotype: IgG1	Calculated MW: 783 aa, 87 kDa	
Immunogen Catalog Number: AG17850	Observed MW: 90-95 kDa	

Applications

Tested Applications:
FC, IF, IHC, WB, ELISA

Species Specificity:
Human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HeLa cells, HEK-293 cells, HepG2 cells, Jurkat cells, K-562 cells, HL-60 cells, THP-1 cells

IHC: human colon cancer tissue, human colon tissue

IF: HepG2 cells,

Background Information

DDX21 protein belongs to DEAD box protein family which is characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD). As a putative RNA helicase, DDX21 unwinds double-stranded RNA, folds single-stranded RNA and is involved in process including ribosomal RNA biogenesis, RNA editing and general transcription. Interaction of DDX21 and c-Jun was reported in ribosomal RNA processing.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

***** 20ul sizes contain 0.1% BSA**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66925-1-Ig (DDX21 antibody) at dilution of 1:50000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control.




Various lysates were subjected to SDS PAGE followed by western blot with 66925-1-Ig (DDX21 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 66925-1-Ig (DDX21 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using 66925-1-Ig (DDX21 antibody) at dilution of 1:100 and CoraLite488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



1×10^6 HepG2 cells were intracellularly stained with 0.2 ug Anti-Human DDX21 (66925-1-Ig, Clone:1B9C8) and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.2 ug Mouse IgG1 Isotype Control (66360-1-Ig, Clone: T1F8D3F10) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).