For Research Use Only

ZFP36 Monoclonal antibody

Catalog Number:66938-1-lg Featured Product 3 Publications




Basic Information	Catalog Number: 66938-1-lg	GenBank Accession Number: BC009693	Purification Method: Protein A purification	
	Size:	GeneID (NCBI):	CloneNo.:	
	150ul , Concentration: 2000 ug/ml by		145E6	
	Nanodrop and 1000 ug/ml by Bradfor		Recommended Dilutions:	
	method using BSA as the standard;	P26651	WB 1:1000-1:6000	
	Source:	Full Name:		
	Mouse	zinc finger protein 36, C3H type, homolog (mouse) Calculated MW:		
	Isotype:			
	IgG2a Immunogen Catalog Number: AG28473			
		326 aa, 34 kDa		
		Observed MW: 34 kDa		
Applications	Tested Applications:	Positive Controls:		
	WB, ELISA	WB : Hep	WB : HepG2 cells, A549 cells, NCI-H1299 cells, LPS	
	Cited Applications: WB, IF, IP	treated U-937 cells, 3T3-L1 cells, pig lung tissue, RAV 264.7 cells		
	Species Specificity:			
	Human, mouse, pig			
Background Informatior	Cited Species: human The expression of many cytokines is	• • • •	y factors that modulate mRNA transport,	
Background Informatior	Cited Species: human The expression of many cytokines is translation, and stability. Much of this mRNAs by proteins that recognize add transcripts. Zfp36 is a mRNA-binding containing mRNAs. It was demonstrat decreased nuclear import and dimini	s regulation occurs by the binding enosine and uridine-rich elements protein involved in post-transcrip ted to physically interact with the shed transcriptional activation mo nRNAs and promoting their degra	y factors that modulate mRNA transport, and stabilizing, or destabilizing, of cytokin s (AREs) in untranslated regions of target tional regulation of AU-rich element (ARE)- p65 subunit of nuclear factor-kB leading to ediated by nuclear factor-kB. It acted by dation, and has a crucial role in the post-	
	Cited Species: human The expression of many cytokines is translation, and stability. Much of this mRNAs by proteins that recognize add transcripts. Zfp36 is a mRNA-binding containing mRNAs. It was demonstrat decreased nuclear import and dimini specifically binding ARE-containing r transcriptional regulation of tumor ne	s regulation occurs by the binding enosine and uridine-rich elements protein involved in post-transcrip ted to physically interact with the shed transcriptional activation mo nRNAs and promoting their degra	and stabilizing, or destabilizing, of cytokin s (AREs) in untranslated regions of target tional regulation of AU-rich element (ARE)- p65 subunit of nuclear factor-kB leading to ediated by nuclear factor-kB. It acted by	
	Cited Species: human The expression of many cytokines is translation, and stability. Much of this mRNAs by proteins that recognize add transcripts. Zfp36 is a mRNA-binding containing mRNAs. It was demonstrated decreased nuclear import and diminis specifically binding ARE-containing re transcriptional regulation of tumor net	s regulation occurs by the binding enosine and uridine-rich elements protein involved in post-transcrip ted to physically interact with the shed transcriptional activation mo nRNAs and promoting their degra- ecrosis factor (TNF).	and stabilizing, or destabilizing, of cytokin s (AREs) in untranslated regions of target tional regulation of AU-rich element (ARE)- p65 subunit of nuclear factor-kB leading to ediated by nuclear factor-kB. It acted by dation, and has a crucial role in the post- Application	
	Cited Species: human The expression of many cytokines is translation, and stability. Much of this mRNAs by proteins that recognize add transcripts. Zfp36 is a mRNA-binding containing mRNAs. It was demonstrat decreased nuclear import and dimini- specifically binding ARE-containing r transcriptional regulation of tumor net Author Pubr Kaiyong Yang 3233	a regulation occurs by the binding enosine and uridine-rich elements protein involved in post-transcrip ted to physically interact with the shed transcriptional activation mo nRNAs and promoting their degra ecrosis factor (TNF).	and stabilizing, or destabilizing, of cytokin s (AREs) in untranslated regions of target tional regulation of AU-rich element (ARE)- p65 subunit of nuclear factor-kB leading to ediated by nuclear factor-kB. It acted by dation, and has a crucial role in the post- Application	
Background Informatior	Cited Species: human The expression of many cytokines is translation, and stability. Much of this mRNAs by proteins that recognize add transcripts. Zfp36 is a mRNA-binding containing mRNAs. It was demonstrat decreased nuclear import and diminis specifically binding ARE-containing r transcriptional regulation of tumor net Author Pubr Kaiyong Yang 3233 Mengge Yin 3954	a regulation occurs by the binding enosine and uridine-rich elements protein involved in post-transcrip ted to physically interact with the shed transcriptional activation me nRNAs and promoting their degra- ecrosis factor (TNF). ned ID Journal 39920 Int Immunopharm	and stabilizing, or destabilizing, of cytokin s (AREs) in untranslated regions of target tional regulation of AU-rich element (ARE)- p65 subunit of nuclear factor-kB leading to ediated by nuclear factor-kB. It acted by dation, and has a crucial role in the post- Application nacol WB	
	Cited Species: human The expression of many cytokines is translation, and stability. Much of this mRNAs by proteins that recognize add transcripts. Zfp36 is a mRNA-binding containing mRNAs. It was demonstrated decreased nuclear import and dimini- specifically binding ARE-containing re transcriptional regulation of tumor net Author Pubr Kaiyong Yang 3233 Mengge Yin 3954	s regulation occurs by the binding enosine and uridine-rich elements protein involved in post-transcrip ted to physically interact with the shed transcriptional activation me nRNAs and promoting their degra- terrosis factor (TNF).	and stabilizing, or destabilizing, of cytokin s (AREs) in untranslated regions of target tional regulation of AU-rich element (ARE)- p65 subunit of nuclear factor-kB leading to ediated by nuclear factor-kB. It acted by dation, and has a crucial role in the post- macol WB WB	
Notable Publications	Cited Species: human The expression of many cytokines is translation, and stability. Much of this mRNAs by proteins that recognize add transcripts. Zfp36 is a mRNA-binding containing mRNAs. It was demonstrated decreased nuclear import and dimini- specifically binding ARE-containing re transcriptional regulation of tumor net Author Pubr Kaiyong Yang 3233 Mengge Yin 3954	er shipment.	and stabilizing, or destabilizing, of cytokin s (AREs) in untranslated regions of target tional regulation of AU-rich element (ARE)- p65 subunit of nuclear factor-kB leading to ediated by nuclear factor-kB. It acted by dation, and has a crucial role in the post- macol WB WB	
	Cited Species: human The expression of many cytokines is translation, and stability. Much of this mRNAs by proteins that recognize add transcripts. Zfp36 is a mRNA-binding containing mRNAs. It was demonstrat decreased nuclear import and dimini: specifically binding ARE-containing r transcriptional regulation of tumor ne Author Pubr Kaiyong Yang 3233 Mengge Yin 3954 Yue Li 3721 Storage: Stora at -20°C. Stable for one year afte Storage Buffer:	s regulation occurs by the binding enosine and uridine-rich elements protein involved in post-transcrip ted to physically interact with the shed transcriptional activation me nRNAs and promoting their degra- scrosis factor (TNF). ned ID Journal 19920 Int Immunopharm 17416 Virus Res 10725 Cell Rep er shipment.	and stabilizing, or destabilizing, of cytokin s (AREs) in untranslated regions of target tional regulation of AU-rich element (ARE)- p65 subunit of nuclear factor-kB leading to ediated by nuclear factor-kB. It acted by dation, and has a crucial role in the post- macol WB WB	

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



HepG2 cells were subjected to SDS PAGE followed by western blot with 66938-1-1g (ZFP36 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.