

For Research Use Only

BLNK Monoclonal antibody

Catalog Number: 66953-1-Ig



Basic Information

Catalog Number: 66953-1-Ig	GenBank Accession Number: BC018906	Purification Method: Protein G purification
Size: 150ul , Concentration: 2400 µg/ml by Nanodrop and 1000 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 29760	CloneNo.: 1A1A1
Source: Mouse	Full Name: B-cell linker	Recommended Dilutions: WB 1:3000-1:8000 IHC 1:250-1:1000 IF 1:200-1:800
Isotype: IgG1	Calculated MW: 50 kDa	
Immunogen Catalog Number: AG28673	Observed MW: 65-70 kDa	

Applications

Tested Applications:

IF, IHC, WB, ELISA

Species Specificity:

Human, pig, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : human spleen tissue, Raji cells, Daudi cells, Ramos cells, pig spleen tissue

IHC : human appendicitis tissue, mouse thymus tissue, rat spleen tissue

IF : Raji cells,

Background Information

BLNK (also known as SLP-65 or BASH) is an important adaptor protein selectively expressed in B-lineage cells. BLNK bridges B cell receptor-associated kinase activation with downstream signaling pathways, thereby affecting various biological functions. BLNK has been shown to be necessary for BCR-mediated Ca²⁺ mobilization, for the activation of mitogen-activated protein kinases such as ERK, JNK, and p38 in a chicken B cell line DT40, and for activation of transcription factors such as NF-AT and NF-κB in human or mouse B cells. BLNK plays a crucial role in pre-BCR-dependent progression of B cell development, BCR-mediated B cell survival, activation, proliferation, and T-independent immune responses.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




human spleen tissue were subjected to SDS PAGE followed by western blot with 66953-1-Ig (BLNK antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human appendicitis tissue slide using 66953-1-Ig (BLNK antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed Raji cells using BLNK antibody (66953-1-Ig, Clone: 1A1A1) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunohistochemical analysis of paraffin-embedded human appendicitis tissue slide using 66953-1-Ig (BLNK antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).