

For Research Use Only

NFATC1 Monoclonal antibody

Catalog Number: 66963-1-Ig

Featured Product

59 Publications



Basic Information

Catalog Number:

66963-1-Ig

Size:

150ul, Concentration: 2000 ug/ml by Nanodrop;

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG13014

GenBank Accession Number:

BC104753

GeneID (NCBI):

4772

UNIPROT ID:

O95644

Full Name:

nuclear factor of activated T-cells, cytoplasmic, calcineurin-dependent 1

Calculated MW:

716aa, 78 kDa; 943aa, 101 kDa

Observed MW:

101 kDa

Purification Method:

Protein G purification

CloneNo.:

1E1B10

Recommended Dilutions:

WB: 1:2000-1:10000

IHC: 1:150-1:600

Applications

Tested Applications:

WB, IHC, ELISA

Cited Applications:

WB, IHC, IF, IP

Species Specificity:

human

Cited Species:

human, mouse, rat, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: Jurkat cells, U2OS cells, Daudi cells, Romas cells, Raji cells, A549 cells, MOLT-4 cells, Ramos cells, Karpas-422 cells

IHC: human urothelial carcinoma tissue, human lymphoma tissue

Background Information

NFATc1(nuclear factor-activated T cells c1) is one of the transcription factors family consisting 5 members which were considered as a regulator of T-cell activation. In immune cell, NFATc1 is located in the cytoplasm in a hyperphosphorylated state and translocates into the nucleus upon cell stimulation via dephosphorylation(PMID:28327458). One paper observed that ectopic expression and nuclear accumulation of NFATc1 in more than 60% of human pancreatic cancers (PMID:25623042). Several alternative splicing transcript variants encoding different isoforms have been identified for this gene. And different isoforms of this protein may regulate different cytokine gene expression. NFATc1 is a main transcriptional activator for cytokine genes in activated lymphoid cells. Also some studies demonstrated that NFATc1 plays the role of a master transcription regulator of osteoclast differentiation (PMID:25489571).

Notable Publications

Author	Pubmed ID	Journal	Application
Huanhuan Xu	35558992	RSC Adv	WB
Beimin Tian	36064833	Int J Oral Sci	IF
Hui-Wen Chiu	34633122	J Bone Miner Res	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66963-1-Ig (NFATC1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human urothelial carcinoma tissue slide using 66963-1-Ig (NFATC1 antibody) at dilution of 1:300 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human urothelial carcinoma tissue slide using 66963-1-Ig (NFATC1 antibody) at dilution of 1:300 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 66963-1-Ig (NFATC1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.