

NFATC1 Monoclonal antibody

Catalog Number: 66963-1-Ig

Featured Product

35 Publications

Basic Information

Catalog Number:

66963-1-Ig

Size:

 150ul, Concentration: 2100 µg/ml by 4772
 Nanodrop and 1000 µg/ml by Bradford
 method using BSA as the standard;

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG13014

GenBank Accession Number:

BC104753

GeneID (NCBI):

4772

UNIPROT ID:

O95644

Full Name:

 nuclear factor of activated T-cells,
 cytoplasmic, calcineurin-dependent 1

Calculated MW:

716aa, 78 kDa; 943aa, 101 kDa

Observed MW:

101 kDa

Purification Method:

Protein G purification

CloneNo.:

1E1B10

Recommended Dilutions:

WB 1:2000-1:10000

IHC 1:150-1:600

Applications

Tested Applications:

IHC, WB, ELISA

Cited Applications:

IF, IHC, IP, WB

Species Specificity:

Human

Cited Species:

human, rat, mouse, pig

**Note-IHC: suggested antigen retrieval with
 TE buffer pH 9.0; (*) Alternatively, antigen
 retrieval may be performed with citrate
 buffer pH 6.0**

Positive Controls:

WB: Jurkat cells, Daudi cells, Romas cells, Raji cells

 IHC: human urothelial carcinoma tissue, human
 lymphoma tissue

Background Information

NFATc1(nuclear factor-activated T cells c1) is one of the transcription factors family consisting 5 members which were considered as a regulator of T-cell activation. In immune cell, NFATc1 is located in the cytoplasm in a hyperphosphorylated state and translocates into the nucleus upon cell stimulation via dephosphorylation(PMID:28327458). One paper observed that ectopic expression and nuclear accumulation of NFATc1 in more than 60% of human pancreatic cancers (PMID:25623042). Several alternative splicing transcript variants encoding different isoforms have been identified for this gene. And different isoforms of this protein may regulate different cytokine gene expression. NFATc1 is a main transcriptional activator for cytokine genes in activated lymphoid cells. Also some studies demonstrated that NFATc1 plays the role of a master transcription regulator of osteoclast differentiation (PMID:25489571).

Notable Publications

Author	Pubmed ID	Journal	Application
Huanhuan Xu	35558992	RSC Adv	WB
Beimin Tian	36064833	Int J Oral Sci	IF
Shuo Ni	32871020	Cell Prolif	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

 T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
 in USA), or 1(312) 455-8498 (outside USA)


 E: proteintech@ptglab.com
 W: ptglab.com

This product is exclusively available under Proteintech
 Group brand and is not available to purchase from any
 other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 66963-1-Ig (NFATC1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human urothelial carcinoma tissue slide using 66963-1-Ig (NFATC1 antibody) at dilution of 1:300 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human urothelial carcinoma tissue slide using 66963-1-Ig (NFATC1 antibody) at dilution of 1:300 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).