

For Research Use Only

CD40 Monoclonal antibody

Catalog Number: 66965-1-Ig 1 Publications



Basic Information

Catalog Number: 66965-1-Ig	GenBank Accession Number: BC012419	Purification Method: Protein A purification
Size: 150ul , Concentration: 1800 ug/ml by Nanodrop and 1000 ug/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 958	CloneNo.: 2A8G5
Source: Mouse	ENSEMBL Gene ID: ENSG00000101017	Recommended Dilutions: WB 1:2000-1:10000 IHC 1:2500-1:10000 IF-P 1:200-1:800 IF/ICC 1:200-1:800
Isotype: IgG1	UNIPROT ID: P25942	
Immunogen Catalog Number: AG28040	Full Name: CD40 molecule, TNF receptor superfamily member 5	
	Calculated MW: 277 aa, 31 kDa	
	Observed MW: 43 kDa	

Applications

Tested Applications: WB, IHC, IF/ICC, IF-P, ELISA	Positive Controls: WB : Raji cells, Daudi cells
Cited Applications: IHC	IHC : human tonsillitis tissue, human lymphoma tissue
Species Specificity: human	IF-P : human tonsillitis tissue,
Cited Species: human	IF/ICC : Ramos cells, Raji cells
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

CD40, also named as TNFRSF5 and Bp50, is a receptor for TNFSF5/CD40LG. It is a member of the TNFR superfamily and is responsible for the efficient activation of cell-mediated and humoral immune responses via activated T cells and B cells respectively. Defects in CD40 lead to hyper-IgM immunodeficiency syndrome type 3 (HIGM3) (PMID: 11675497). This is an autosomal recessive disorder that includes the inability of B cells to undergo isotype switching, a key step in the final differentiation of the humoral immune response, and an inability to mount an antibody-specific immune response.

Notable Publications

Author	Pubmed ID	Journal	Application
Beiyuan Hu	36304515	J Clin Transl Hepatol	IHC

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Raji cells were subjected to SDS PAGE followed by western blot with 66965-1-Ig (CD40 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 66965-1-Ig (CD40 antibody) at dilution of 1:5000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 66965-1-Ig (CD40 antibody) at dilution of 1:5000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed Ramos cells using CD40 antibody (66965-1-Ig, Clone: 2A8G5) at dilution of 1:400 and Multi-rAb CoraLite® Plus 488-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (RGAM002).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using CD40 antibody (66965-1-Ig, Clone: 2A8G5) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using CD40 antibody (66965-1-Ig, Clone: 2A8G5) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).