

For Research Use Only

TRPV1 Monoclonal antibody

Catalog Number: 66983-1-Ig **47 Publications**



Basic Information

Catalog Number: 66983-1-Ig	GenBank Accession Number: BC132820	Purification Method: Protein A purification
Size: 150ul , Concentration: 1000 ug/ml by Nanodrop;	GeneID (NCBI): 7442	CloneNo.: 1A3C9
Source: Mouse	UNIPROT ID: Q8NER1	Recommended Dilutions: WB 1:1000-1:6000
Isotype: IgG1	Full Name: transient receptor potential cation channel, subfamily V, member 1	
Immunogen Catalog Number: AG18683	Calculated MW: 839 aa, 95 kDa	
	Observed MW: 105 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : HepG2 cells, MDA-MB-468 cells, T-47D cells, HEK-293 cells, Y79 cells, HUVEC cells
Cited Applications: WB, ColP	
Species Specificity: Human	
Cited Species: human, sheep	

Background Information

TRPV1, also named as VR1 or OTRPC1, belongs to the transient receptor potential (TRP) family and TRPV subfamily (PMID: 17579562). TRPV1 is a ligand-activated non-selective calcium permeant cation channel expressed predominantly by sensory neurons (PMID: 10821274). It is involved in detection of noxious chemical and thermal stimuli. TRPV1 plays an important role in inflammatory thermal hyperalgesia (PMID: 10764638; 10821274).

Notable Publications

Author	Pubmed ID	Journal	Application
Zhongyang Lv	36137413	EBioMedicine	WB
Sampurna Chakrabarti	34544757	eNeuro	
Dingding Liu	35350752	Front Pharmacol	WB

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



HepG2 cells were subjected to SDS PAGE followed by western blot with 66983-1-Ig (TRPV1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.