

For Research Use Only

TGM1 Monoclonal antibody

Catalog Number: 67004-1-Ig



Basic Information

Catalog Number: 67004-1-Ig	GenBank Accession Number: BC034699	Purification Method: Protein A purification
Size: 150ul , Concentration: 1800 µg/ml by Nanodrop and 1000 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 7051	CloneNo.: 3D9C11
Source: Mouse	Full Name: transglutaminase 1 (K polypeptide epidermal type I, protein-glutamine- gamma-glutamyltransferase)	Recommended Dilutions: WB 1:1000-1:6000 IHC 1:500-1:2000
Isotype: IgG2b	Calculated MW: 817 aa, 90 kDa	
Immunogen Catalog Number: AG3905	Observed MW: 90 kDa	

Applications

Tested Applications:

IHC, WB, ELISA

Species Specificity:

Human

**Note-IHC: suggested antigen retrieval with
TE buffer pH 9.0; (*) Alternatively, antigen
retrieval may be performed with citrate
buffer pH 6.0**

Positive Controls:

WB : human saliva tissue,

IHC : human tonsillitis tissue, human cervical cancer
tissue

Background Information

Transglutaminase 1 is an epidermal enzyme that is expressed during the terminal differentiation of keratinized squamous epithelium to form cornified cell envelope in differentiated keratinocytes. Mutation of TGM1 has been linked to various epidermal diseases. This antibody recognizes the endogenous 90 kDa TGM1 proteins.

Storage

Storage:

Store at -20°C.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

***** 20ul sizes contain 0.1% BSA**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

**This product is exclusively available under Proteintech
Group brand and is not available to purchase from any
other manufacturer.**

Selected Validation Data



human saliva were subjected to SDS PAGE followed by western blot with 67004-1-Ig (TGM1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 67004-1-Ig (TGM1 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).