## For Research Use Only

## MAP2 Monoclonal antibody, PBS Only

Catalog Number:67015-1-PBS



**Basic Information** 

Catalog Number:

GenBank Accession Number:

**Purification Method:** Protein A purification

67015-1-PBS

BC038857 GeneID (NCBI):

Size:

100ug, Concentration: 1mg/ml by 4133 CloneNo.: 1C3E6

Nanodrop;

**UNIPROT ID:** 

Source:

P11137 Full Name:

Mouse Isotype:

microtubule-associated protein 2

lgG2b

Calculated MW:

Immunogen Catalog Number:

200 kDa

AG11349

**Applications** 

**Tested Applications:** 

IHC, IF-P, IF-Fro, FC (Intra), Indirect ELISA

Species Specificity:

human, mouse, rat

**Background Information** 

MAP2 (microtubule-associated protein 2) is a cytoskeleton protein abundant in brain and has important role in neuronal morphogenesis. Multiple high molecular weight (MW) and low molecular weight (MW) MAP2 isoforms are expressed within axons, dendrites, and cell bodies. The expression of MAP2 is regulated in both a tissue- and developmentally specific manner. MAP2 antibodies have been widely used to mark the neuron or dendrite formation.

Storage

Storage:

Store at -80°C.

Storage Buffer:


PBS only, pH7.3

in USA), or 1(312) 455-8498 (outside USA)


## Selected Validation Data




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 67015-1-Ig (MAP2 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67015-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed rat brain tissue using MAP2 antibody (67015-1-lg, Clone: 1C3E6) at dilution of 1:400 and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+I). This data was developed using the same antibody clone with 67015-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using MAP2 antibody (67015-1-Ig, Clone: 1C3E6) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 67015-1-PBS in a different storage buffer formulation.




1X10^6 Neuro-2a cells were intracellularly stained with 0.4 ug Anti-Human MAP2 (67015-1-Ig, Clone:1C3E6) and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG2b Isotype Control (MPC-11) (65128-1-Ig, Clone: MPC-11) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 67015-1-PBS in a




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 67015-1-Ig (MAP2 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67015-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 67015-1-lg (MAP2 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67015-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 67015-1-lg (MAP2 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67015-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using MAP2 antibody (67015-1-1g, Clone: 1C3E6) at dilution of 1:400 and CoraLite® 488-Conjugated Affini Pure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 67015-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed frozen OCT-embedded mouse brain tissue using MAP2 antibody (67015-1-lg, Clone: 1C3E6) at dilution of 1:800 and Coralite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) (SA00013-1). This data was developed using the same antibody clone with 67015-1-PBS in a different storage buffer formulation.