### For Research Use Only

# TIMELESS Monoclonal antibody

Catalog Number:67022-1-lg Featured Product

1 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number:

67022-1-lg BC050557 GeneID (NCBI): Size:

150ul, Concentration: 2200 ug/ml by 8914 Nanodrop and 1000 ug/ml by Bradford<sub>UNIPROT ID:</sub> method using BSA as the standard; Q9UNS1

Source: Full Name:

Mouse timeless homolog (Drosophila)

Isotype: Calculated MW: lgG1 139 kDa Immunogen Catalog Number: Observed MW: AG6283 140 kDa

**Purification Method:** 

Protein A purification

CloneNo.: 1B6G3

Recommended Dilutions: WB 1:5000-1:50000

**Applications** 

**Tested Applications:** WB, FC (Intra), ELISA **Cited Applications:** 

WB

Species Specificity: human, mouse, rat Cited Species: human, canine

#### Positive Controls:

WB: HeLa cells, U2OS cells, A549 cells, JAR cells, Jurkat cells, HSC-T6 cells, NIH/3T3 cells, HEK-293

cells, K-562 cells

# **Background Information**

 ${\bf TIMELESS1} \ ({\bf TIM1}) \ gene \ encodes \ protein \ timeless \ homolog \ belonging \ to \ timeless \ family. \ TIM1 \ is \ localized \ in$ nucleus and regulated by cell cycle. It expresses high levels in S, G2 and M phases, and low in G0 and G1 phases. TIM1 is involved in the circadian rhythm autoregulatory loop and is a crucial component of the circadian core oscillator, which includes the CRY proteins, CLOCK, NPAS2, etc. It is also involved in cell survival after DNA damage or replication stress and reported to play important role in epithelial cell morphogenesis and formation of branching tubules.

#### **Notable Publications**

| Author      | Pubmed ID | Journal         | Application |
|-------------|-----------|-----------------|-------------|
| Houqing Yin | 37433364  | Eur J Pharmacol | WB          |

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 67022-1-1g (TIMELESS antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Lamin B1 Monoclonal antibody (HRP-66095) as loading control.



1X10^6 HEK-293 cells were intracellularly stained with 0.4 ug Anti-Human TIMELESS (67022-1-Ig, Clone:1B6G3) and Coralite®488-Conjugated Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).