

For Research Use Only

PELP1 Monoclonal antibody

Catalog Number: 67050-1-Ig **2 Publications**



Basic Information

Catalog Number: 67050-1-Ig	GenBank Accession Number: BC069058	Purification Method: Protein A purification
Size: 150ul, Concentration: 1000 ug/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 27043	CloneNo.: 1B11E4
Source: Mouse	UNIPROT ID: Q8IZL8	Recommended Dilutions: WB 1:2000-1:8000 IHC 1:500-1:2000 IF-P 1:200-1:800 IF-Fro 1:200-1:800
Isotype: IgG2b	Full Name: proline, glutamate and leucine rich protein 1	
Immunogen Catalog Number: AG25729	Calculated MW: 1130 aa, 120 kDa	
	Observed MW: 160 kDa	

Applications

Tested Applications: WB, IHC, IF-P, IF-Fro, FC (Intra), ELISA	Positive Controls: WB : HeLa cells, T-47D cells, Jurkat cells, HEK-293 cells, MCF-7 cells IHC : human cervical cancer tissue, human breast cancer tissue IF-P : human breast cancer tissue, IF-Fro : mouse breast cancer,
Cited Applications: WB, IHC, IF	
Species Specificity: human	
Cited Species: mouse	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

PELP1 was first identified as a 160 kDa protein in a screen for Src homology 2 (SH2) domain-binding proteins. PELP1 is overexpressed in 60-80% of breast tumors and plays important roles in both ER genomic and non-genomic signaling. In vivo, PELP1 subcellular localization is primarily nuclear in normal breast tissue, but it is localized to the cytoplasm in about 40% of invasive breast tumors. In the nucleus, PELP1 interacts with a number of transcription factors. The proto-oncogenic functions of PELP1 involve different cellular processes including epigenetic modifications leading to ER transactivation and breast cancer progression. Furthermore, PELP1 activates kinase cascades in the cytoplasm such as MAPK activation via c-Src and PI3K signaling.

Notable Publications

Author	Pubmed ID	Journal	Application
Jinzhao Ning	38664304	Mol Biotechnol	WB,IHC,IF
Kai-Xiang He	37290329	Int Immunopharmacol	WB,IHC

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67050-1-Ig (PELP1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human cervical cancer tissue slide using 67050-1-Ig (PELP1 antibody) at dilution of 1:1000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human cervical cancer tissue slide using 67050-1-Ig (PELP1 antibody) at dilution of 1:1000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed human breast cancer tissue using PELP1 antibody (67050-1-Ig, Clone: 1B11E4) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



1X10⁶ MCF-7 cells were intracellularly stained with 0.4 ug Anti-Human PELP1 (67050-1-Ig, Clone:1B11E4) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG2b Isotype Control (66360-3-Ig, Clone: K11B8C4B5) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Immunofluorescent analysis of (4% PFA) fixed human breast cancer tissue using PELP1 antibody (67050-1-Ig, Clone: 1B11E4) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed frozen OCT-embedded mouse breast cancer using PELP1 antibody (67050-1-Ig, Clone: 1B11E4) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1).