For Research Use Only

PELP1 Monoclonal antibody, PBS Only

Catalog Number: 67050-1-PBS



Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method:

67050-1-PBS

BC069058 GeneID (NCBI): Protein A purification

27043

CloneNo.:

1B11E4

100ug, Concentration: 1mg/ml by Nanodrop:

UNIPROT ID:

Q8IZL8

Full Name:

Mouse Isotype: lgG2b

proline, glutamate and leucine rich

protein 1

Immunogen Catalog Number:

Calculated MW:

AG25729

1130 aa, 120 kDa

Observed MW:

160 kDa

Applications

Tested Applications:

WB, IHC, IF-P, IF-Fro, FC (Intra), Indirect ELISA

Species Specificity:

Background Information


PELP1 was first identified as a 160 kDa protein in a screen for Src homology 2 (SH2) domain-binding proteins. PELP1 is overexpressed in 60-80% of breast tumors and plays important roles in both ER genomic and non-genomic signaling. In vivo, PELP1 subcellular localization is primarily nuclear in normal breast tissue, but it is localized to the cytoplasm in about 40% of invasive breast tumors. In the nucleus, PELP1 interacts with a number of transcription factors. The proto-oncogenic functions of PELP1 involve different cellular processes including epigenetic modifications leading to ER transactivation and breast cancer progression. Furthermore, PELP1 activates kinase cascades in the cytoplasm such as MAPK activation via c-Src and PI3K signaling.

Storage


Storage: Store at -80°C. Storage Buffer: PBS only, pH7.3

in USA), or 1(312) 455-8498 (outside USA)


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67050-1-lg (PELP1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67050-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human cervical cancer tissue slide using 67050-1-Ig (PELP1 antibody) at dilution of 1:1000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67050-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human cervical cancer tissue slide using 67050-1-lg (PELP1 antibody) at dilution of 1:1000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67050-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed human breast cancer tissue using PELP1 antibody (67050-1-Ig, Clone: 1B11E4) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 67050-1-PBS in a different storage buffer formulation.



1X10^6 MCF-7 cells were intracellularly stained with 0.4 ug Anti-Human PELP1 (67050-1-lg, Clone:1B11E4) and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG2b Isotype Control (66360-3-lg, Clone: K11B8C4B5) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). This data was developed using the same antibody clone with 67050-1-PBS



Immunofluorescent analysis of (4% PFA) fixed human breast cancer tissue using PELP1 antibody (67050-1-Ig, Clone: 1B11E4) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 67050-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed frozen OCT-embedded mouse breast cancer using PELP1 antibody (67050-1-Ig, Clone: 1B11E4) at dilution of 1:400 and CoralLite@488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1). This data was developed using the same antibody clone with 67050-1-PBS in a different storage buffer formulation