

For Research Use Only

REDD1 Monoclonal antibody

Catalog Number: 67059-1-Ig

Featured Product

2 Publications



Basic Information

Catalog Number: 67059-1-Ig	GenBank Accession Number: BC007714	Purification Method: Protein A purification
Size: 150ul, Concentration: 1600 ug/ml by Nanodrop and 1000 ug/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 54541	CloneNo.: 3A2C10
Source: Mouse	UNIPROT ID: Q9NX09	Recommended Dilutions: WB 1:1000-1:8000
Isotype: IgG1	Full Name: DNA-damage-inducible transcript 4	
Immunogen Catalog Number: AG0965	Calculated MW: 25 kDa	
	Observed MW: 32-35 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : PC-3 cells, HeLa cells, LNCaP cells, HepG2 cells, A549 cells, K-562 cells
Cited Applications: WB	
Species Specificity: Human	
Cited Species: human	

Background Information

REDD1, also named as RTP801 and DDIT4, belongs to the DDIT4 family. REDD1 promotes neuronal cell death. It is a novel transcriptional target of p53 implicated ROS in the p53-dependent DNA damage response. REDD1 controlled cell growth under energy stress, as an essential regulator of TOR activity through the TSC1/2 complex. REDD-1 expression has also been linked to apoptosis, A β toxicity and the pathogenesis of ischemic diseases. As an HIF-1-responsive gene, REDD-1 exhibits strong hypoxia-dependent upregulation in ischemic cells of neuronal origin [PMID: 19996311]. In response to stress due to DNA damage and glucocorticoid treatment, REDD-1 is upregulated at the transcriptional level [PMID: 21733849]. REDD-1 negatively regulates the mammalian target of Rapamycin, a serine/threonine kinase often referred to as mTOR [PMID: 22951983]. It is crucial in the coupling of extra- and intracellular cues to mTOR regulation. The absence of REDD-1 is associated with the development of retinopathy, a major cause of blindness [PMID: 22304497]. REDD1 is a new host defense factor, and chemical activation of REDD1 expression represents a potent antiviral intervention strategy [PMID: 21909097]. The calculated molecular weight of REDD1 is 25 kDa. Because of multiple lysines in the proteins, REDD1 often migrates around 35 kDa on Western blot [PMID: 19221489].

Notable Publications

Author	Pubmed ID	Journal	Application
Guodan Zeng	34102031	FEBS Open Bio	WB
Yi Zhang	37671155	Front Immunol	WB

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67059-1-Ig (REDD1 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.