For Research Use Only

ATF5 Monoclonal antibody

Catalog Number:67066-1-lg 3 Publications




Basic Information	Catalog Number: 67066-1-lg	GenBank Accession Num BC005174	nber:	Purification Method: Protein A purification
	Size:	GeneID (NCBI):		CloneNo.:
	150ul , Concentration: 2300 ug/ml by			1E5B6
	Nanodrop and 1000 ug/ml by Bradfor method using BSA as the standard;	^d UNIPROT ID: Q9Y2D1		Recommended Dilutions: WB 1:2000-1:16000
	Source: Mouse	Full Name: activating transcription	factor 5	
	lsotype: lgG2b	Calculated MW: 31 kDa		
	Immunogen Catalog Number: AG7214	Observed MW: 30-35 kDa		
Applications	Tested Applications: WB, ELISA	Ρ	Positive Cont	rols:
	WB, ELISA Cited Applications: WB	c	ells, HeLa cells, HEK-293 cells, SH-SY5Y ells, HT-29 cells, MCF-7 cells, MDA-MB- BR-3 cells, T-47D cells	
	Species Specificity: Human			
	Cited Species: human, chicken, mouse			
	namer, circler, mode			
Background Information	ATF5, also named as Activating trans ATF5 is Widely expressed with highe accumulated in the pericentriolar ma During late telophase and cytokinesi as a transcription factor that either sti regulatory elements such as cAMP res response element (ARE) (consensus: 5	r expression levels in live terial (PCM) during G1 to s, it translocates from the imulates or represses gen sponse element (CRE) (con 5'-C[CT]TCT[CT]CCTT[AT] ponoters. Critically ATF5 is	er. ATF5 is ac M phase via centrosome he transcriptio msensus: 5'-G]-3') but also s involved, o	iTGACGT[AC][AG]-3'), ATF5-specific the amino acid response element (AARE ften in a cell type-dependent manner, in
	ATF5, also named as Activating trans ATF5 is Widely expressed with highe accumulated in the pericentriolar ma During late telophase and cytokinesi: as a transcription factor that either sti regulatory elements such as cAMP re- response element (ARE) (consensus: 5 present in many viral and cellular pro cell survival, proliferation, and differ PubMed:20654631).	r expression levels in live terial (PCM) during G1 to s, it translocates from the imulates or represses gen sponse element (CRE) (con 5'-C[CT]TCT[CT]CCTT[AT] ponoters. Critically ATF5 is	er. ATF5 is ac M phase via centrosome te transcriptionsensus: 5'-G [-3') but also s involved, o 5550, PubMed	tively transported to the centrosome and a microtubule-dependent mechanism. to the midbody (PMID: 26213385). ATF5 on through binding of different DNA GACGT[AC][AG]-3'), ATF5-specific the amino acid response element (AARE fften in a cell type-dependent manner, in
	ATF5, also named as Activating trans ATF5 is Widely expressed with higher accumulated in the pericentriolar ma During late telophase and cytokinesia as a transcription factor that either stir regulatory elements such as cAMP re- response element (ARE) (consensus: 5 present in many viral and cellular pro- cell survival, proliferation, and differe PubMed:20654631).	r expression levels in live terial (PCM) during G1 to s, it translocates from the imulates or represses gen sponse element (CRE) (con 5'-C[CT]TCT[CT]CCT[AT] omoters. Critically ATF5 is entiation (PubMed:10373)	er. ATF5 is ac M phase via centrosome te transcripti insensus: 5'-G]-3') but also is involved, o 5550, PubMed	trively transported to the centrosome and a microtubule-dependent mechanism. to the midbody (PMID: 26213385) . ATF5 on through binding of different DNA iTGACGT[AC][AG]-3'), ATF5-specific the amino acid response element (AARE ften in a cell type-dependent manner, in :15358120, PubMed:21212266,
	ATF5, also named as Activating trans ATF5 is Widely expressed with highe accumulated in the pericentriolar ma During late telophase and cytokinesis as a transcription factor that either sti regulatory elements such as cAMP re- response element (ARE) (consensus: 9 present in many viral and cellular pro- cell survival, proliferation, and differe PubMed:20654631). Author Pul Kosei Yamashita 362	r expression levels in live terial (PCM) during G1 to s, it translocates from the imulates or represses gen sponse element (CRE) (con 5'-C[CT]TCT[CT]CCTT[AT] omoters. Critically ATF5 is entiation (PubMed:10373)	er. ATF5 is ac M phase via centrosome le transcriptionsensus: 5'-G]-3') but also s involved, o (550, PubMed	trively transported to the centrosome and a microtubule-dependent mechanism. to the midbody (PMID: 26213385) . ATF5 on through binding of different DNA iTGACGT[AC][AG]-3'), ATF5-specific the amino acid response element (AARE fiten in a cell type-dependent manner, in 15358120, PubMed:21212266, Application
	ATF5, also named as Activating trans ATF5 is Widely expressed with higher accumulated in the pericentriolar ma During late telophase and cytokinesia as a transcription factor that either stir regulatory elements such as cAMP res- response element (ARE) (consensus: 5 present in many viral and cellular pro- cell survival, proliferation, and differ PubMed:20654631). Author Pul Kosei Yamashita 362 Meiyu Cheng 353	r expression levels in live terial (PCM) during G1 to s, it translocates from the imulates or represses gen sponse element (CRE) (con 5'-C[CT]TCT[CT]CCTT[AT] omoters. Critically ATF5 is entiation (PubMed:10373) bmed ID Journal 217717 FEBS Op	er. ATF5 is ac M phase via centrosome te transcriptionsensus: 5'-G]-3') but also s involved, o 5550, PubMed	tively transported to the centrosome and a microtubule-dependent mechanism. to the midbody (PMID: 26213385) . ATF5 on through binding of different DNA iTGACGT[AC][AG]-3'), ATF5-specific the amino acid response element (AARE fiten in a cell type-dependent manner, in 15358120, PubMed:21212266, Application WB
Background Information Notable Publications	ATF5, also named as Activating trans ATF5 is Widely expressed with higher accumulated in the pericentriolar ma During late telophase and cytokinesia as a transcription factor that either stir regulatory elements such as cAMP res- response element (ARE) (consensus: 5 present in many viral and cellular pro- cell survival, proliferation, and differ PubMed:20654631). Author Pul Kosei Yamashita 362 Meiyu Cheng 353	r expression levels in live terial (PCM) during G1 to s, it translocates from the imulates or represses gen sponse element (CRE) (con 5'-C[CT]TCT[CT]CCT[AT] omoters. Critically ATF5 i: entiation (PubMed:10373) bmed ID Journal 217717 FEBS Op 163244 Int J Mo 748667 Toxicol er shipment. % glycerol pH 7.3.	er. ATF5 is ac M phase via centrosome te transcriptionsensus: 5'-G]-3') but also s involved, o 5550, PubMed	trively transported to the centrosome and a microtubule-dependent mechanism. to the midbody (PMID: 26213385) . ATF5 on through binding of different DNA iTGACGT[AC][AG]-3'), ATF5-specific the amino acid response element (AARE fiten in a cell type-dependent manner, in hts558120, PubMed:21212266, Application WB WB

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67066-1-1g (ATF5 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.