

For Research Use Only

TFAP2A/AP-2 Monoclonal antibody

Catalog Number: 67076-1-Ig **2 Publications**



Basic Information

Catalog Number: 67076-1-Ig	GenBank Accession Number: BC017754	Purification Method: Protein A purification
Size: 150ul, Concentration: 2100 ug/ml by Nanodrop and 900 ug/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 7020	CloneNo.: 1B12F11
Source: Mouse	UNIPROT ID: P05549	Recommended Dilutions: WB 1:2000-1:10000 IHC 1:250-1:1000
Isotype: IgG1	Full Name: transcription factor AP-2 alpha (activating enhancer binding protein 2 alpha)	
Immunogen Catalog Number: AG4112	Calculated MW: 431 aa, 47 kDa	
	Observed MW: 47 kDa	

Applications

Tested Applications: WB, IHC, FC (Intra), ELISA	Positive Controls: WB : LO2 cells, HeLa cells, HSC-T6 cells, MCF-7 cells, HepG2 cells IHC : human breast cancer tissue,
Cited Applications: IHC	
Species Specificity: human, mouse, rat	
Cited Species: human	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

The activator protein-2 (AP-2) family of transcription factors comprises five 52-kDa isoforms (AP-2 α , AP-2 β , AP-2 γ , AP-2 δ , and AP-2 ϵ), which share a common structure: a proline/glutamine-rich transactivation domain in the N-terminal region and a helix-span-helix domain in the C-terminal region, which mediates dimerization and site-specific DNA binding. Depending on the cellular context, the AP-2 transcription factors are individually associated either with cell differentiation and development or with cancer progression/regression. [PMID:21966377] TFAP2A (AP-2-alpha) is the only AP-2 protein required for early morphogenesis of the lens vesicle. Together with the CITED2 coactivator, stimulates the PITX2 P1 promoter transcription activation [PMID:11694877]

Notable Publications

Author	Pubmed ID	Journal	Application
Junyuan Yang	37083077	Oncol Rep	IHC
Feng Zhang	36911415	Front Genet	IHC

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



L02 cells were subjected to SDS PAGE followed by western blot with 67076-1-Ig (TFAP2A/AP-2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 67076-1-Ig (TFAP2A/AP-2 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1X10⁶ A431 cells were intracellularly stained with 0.4 ug Anti-Human TFAP2A/AP-2 (67076-1-Ig, Clone:1B12F11) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG1 Isotype Control (MOPC-21) (65124-1-Ig, Clone: MOPC-21) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).