## For Research Use Only

## ATG9A Monoclonal antibody

Catalog Number:67096-1-lg 5 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number: Purification Method: 67096-1-lg BC065534 Protein G purification

Size: GeneID (NCBI): CloneNo.: 150ul , Concentration: 2500 µg/ml by 79065 1D12E11

Nanodrop and 1000 µg/ml by Bradford Full Name: Recommended Dilutions: method using BSA as the standard; ATC9 autophagy related 9 homolog A WB 1:2000-1:10000

ATG9 autophagy related 9 homolog A WB 1:2000-1:10000
Source: (S. cerevisiae) IHC 1:300-1:1200
Mouse Calculated MW: IF 1:50-1:500

Isotype:837 aa, 94 kDaIgG1Observed MW:Immunogen Catalog Number:94 kDa

AG24278

**Applications** 

Tested Applications: Positive Controls:

IF, IHC, WB, ELISA WB: HEK-293 cells, HepG2 cells, Jurkat cells

Cited Applications: IHC : human thyroid cancer tissue,

Species Specificity: IF: HepG2 cells,

Human
Cited Species:
human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

**Background Information** 

ATG9A is the only transmembrane ATG protein essential for autophagy. It plays a key role in the organization of the preautophagosomal structure/phagophore assembly site (PAS). It has been reported that ATG9A expression is increased in oral squamous cell carcinoma and breast cancers. The inhibition of ATG9A can lead to an inhibition of cancer cell proliferation and invasion.(PMID: 29437695, 29568063)

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Zhiheng Zhang	38228036	Drug Resist Updat	WB
Laurence Lecordier	38041817	Cell Rep	IF
Xuezhao Feng	37957156	Nat Commun	WB,IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 67096-1-1g (ATG9A antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using 67096-1-1g (ATG9A antibody) at dilution of 1:100 and CoraLite488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunohistochemical analysis of paraffinembedded human thyroid cancer tissue slide using 67096-1-Ig (ATG9A antibody) at dilution of 1:600 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).