For Research Use Only

ATG9A Monoclonal antibody

Catalog Number:67096-1-lg 5 Publications




Basic Information	Catalog Number: 67096-1-lg	GenBank Accession Number: BC065534	Purification Method: Protein G purification	
	Size: 150ul , Concentration: 1000 ug/ml by Nanodrop; Source: Mouse	GeneID (NCBI):	CloneNo.:	
		/ 79065	1D12E11	
		UNIPROT ID: Q7Z3C6	Recommended Dilutions: WB 1:2000-1:10000	
		Full Name:	IHC 1:300-1:1200	
	lsotype: lgG1	ATG9 autophagy related 9 homolog A IF/ICC 1:50-1:500 (S. cerevisiae)		
	Immunogen Catalog Number: AG24278	Calculated MW: 837 aa, 94 kDa		
		Observed MW: 94 kDa		
Applications	Tested Applications:	Positive Controls:		
	WB, IHC, IF/ICC, ELISA	WB : HEK-293 cells, HepG2 cells, Jurkat cells		
	Cited Applications: WB, IF	IHC : human thyroid cancer tissue,		
	Species Specificity: Human	IF/ICC :	IF/ICC : HepG2 cells,	
	Cited Species: human, mouse Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0			
Background Informatior	retrieval may be performed w buffer pH 6.0 ATG9A is the only transmembrane A preautophagosomal structure/phago	With citrate TG protein essential for autophag phore assembly site (PAS). It has noma and breast cancers. The inh	been reported that ATG9A expression is	
	retrieval may be performed w buffer pH 6.0 ATG9A is the only transmembrane A preautophagosomal structure/phago increased in oral squamous cell carci cancer cell proliferation and invasion	With citrate TG protein essential for autophag phore assembly site (PAS). It has noma and breast cancers. The inh	y. It plays a key role in the organization of t been reported that ATG9A expression is ibition of ATG9A can lead to an inhibition o Application	
	retrieval may be performed w buffer pH 6.0 ATG9A is the only transmembrane A ^T preautophagosomal structure/phago increased in oral squamous cell carci cancer cell proliferation and invasion Author Put	Fith citrate TG protein essential for autophag phore assembly site (PAS). It has noma and breast cancers. The inh n.(PMID: 29437695, 29568063)	been reported that ATG9A expression is ibition of ATG9A can lead to an inhibition of Application	
Background Informatior Notable Publications	retrieval may be performed w buffer pH 6.0 ATG9A is the only transmembrane A' preautophagosomal structure/phago increased in oral squamous cell carci cancer cell proliferation and invasion Author Put Zhiheng Zhang 382	TG protein essential for autophag phore assembly site (PAS). It has noma and breast cancers. The inh n.(PMID: 29437695, 29568063) pmed ID Journal	been reported that ATG9A expression is ibition of ATG9A can lead to an inhibition of Application	
	retrieval may be performed w buffer pH 6.0 ATG9A is the only transmembrane A preautophagosomal structure/phago increased in oral squamous cell carci cancer cell proliferation and invasion Author Put Zhiheng Zhang 382 Laurence Lecordier 380	TG protein essential for autophag phore assembly site (PAS). It has noma and breast cancers. The inh n.(PMID: 29437695, 29568063) pmed ID Journal 228036 Drug Resist Upd	been reported that ATG9A expression is ibition of ATG9A can lead to an inhibition of Application at WB	
	retrieval may be performed w buffer pH 6.0 ATG9A is the only transmembrane A preautophagosomal structure/phago increased in oral squamous cell carci cancer cell proliferation and invasion Author Put Zhiheng Zhang 382 Laurence Lecordier 380	TG protein essential for autophag phore assembly site (PAS). It has noma and breast cancers. The inh n.(PMID: 29437695, 29568063) omed ID Journal 228036 Drug Resist Upd 041817 Cell Rep 057156 Nat Commun er shipment.	been reported that ATG9A expression is ibition of ATG9A can lead to an inhibition of Application at WB IF	


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67096-1-1g (ATG9A antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using 67096-1-1g (ATG9A antibody) at dilution of 1:100 and CoraLite488-Conjugated Goat Anti-Mouse IgG(H+L).



Immunohistochemical analysis of paraffinembedded human thyroid cancer tissue slide using 67096-1-1g (ATG9A antibody) at dilution of 1:600 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).