

HOXA7 Monoclonal antibody

Catalog Number: 67112-1-Ig

Featured Product

3 Publications

Basic Information

Catalog Number:

67112-1-Ig

Size:

 150ul, Concentration: 2500 ug/ml by
 Nanodrop and 1000 ug/ml by Bradford
 method using BSA as the standard;

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG19025

GenBank Accession Number:

BC148692

GeneID (NCBI):

3204

UNIPROT ID:

P31268

Full Name:

homeobox A7

Calculated MW:

230 aa, 25 kDa

Observed MW:

20 kDa

Purification Method:

Protein G purification

CloneNo.:

2A7H6

Recommended Dilutions:

WB 1:2000-1:10000

IF/ICC 1:50-1:500

Applications

Tested Applications:

WB, IF/ICC, ELISA

Cited Applications:

WB, IHC, IF

Species Specificity:

human

Cited Species:

human, mouse

Positive Controls:

 WB : A431 cells, HCT 116 cells, HepG2 cells, HL-60
 cells, HT-29 cells, K-562 cells, SMMC-7721 cells, THP-1
 cells, ACHN cells, SiHa cells, NCI-H1299 cells, HEK-293
 cells, RKO cells

IF/ICC : HepG2 cells,

Background Information

HOX genes play a fundamental role in the development of the vertebrate central nervous system, heart, axial skeleton, limbs, gut, urogenital tract and external genitalia. The homeobox gene Hoxa-1 is transcriptionally regulated by retinoic acid (RA) and encodes a transcription factor, which has been shown to play important roles in cell differentiation and embryogenesis. Hoxa-1 is also expressed in cancers, such as mammary tumors, though it is not expressed in normal gland or in precancerous mammary tissues. At embryonic stages, Hoxa-2 is expressed in the mesenchyme and epithelial cells of palate, however its expression is restricted to the tips of the growing palatal shelves. Hoxa-2 protein is predominantly expressed in the nuclei of cells in the ventral mantle region of the developing embryo. In the developing and adult mouse spinal cord, Hoxa-2 protein may contribute to dorsal-ventral patterning and/or to the specification of neuronal phenotype. Hoxa-7 functions as a potent transcriptional repressor and its action as such requires several domains, including both activator and repressor regions. Hoxa-7 is expressed in the fetal liver, lung, skeletal muscle, kidney, pancreas and placenta.

Notable Publications

Author	Pubmed ID	Journal	Application
Jun-Pu Zha	32650806	J Orthop Surg Res	WB
Jun Li	34978042	Reprod Sci	WB
Anqi Feng	37248653	Cancer Sci	WB, IHC, IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

 T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
 in USA), or 1(312) 455-8498 (outside USA)


 E: proteintech@ptglab.com
 W: ptglab.com

This product is exclusively available under Proteintech
 Group brand and is not available to purchase from any
 other manufacturer.

Selected Validation Data



A431 cells were subjected to SDS PAGE followed by western blot with 67112-1-Ig (HOXA7 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using 67112-1-Ig (HOXA7 antibody) at dilution of 1:100 and CoraLite488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).