

For Research Use Only

USF1 Monoclonal antibody

Catalog Number: 67114-1-Ig



Basic Information

Catalog Number: 67114-1-Ig	GenBank Accession Number: BC035505	Purification Method: Protein G purification
Size: 150ul , Concentration: 1000 ug/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 7391	CloneNo.: 1C8F1
Source: Mouse	UNIPROT ID: P22415	Recommended Dilutions: WB 1:2000-1:10000
Isotype: IgG1	Full Name: upstream transcription factor 1	
Immunogen Catalog Number: AG17785	Calculated MW: 34 kDa	
	Observed MW: 38-43 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : HeLa cells, HEK-293 cells, HT-29 cells, K-562 cells, HSC-T6 cells, NIH/3T3 cells, Jurkat cells, PC-12 cells, 4T1 cells
Species Specificity: Human, mouse, rat	

Background Information

Upstream stimulatory factor 1 (USF1), also name Class B basic helix-loop-helix protein 11. It is a major late transcription factor 1. USF1 is a member of the basic helix-loop-helix leucine zipper family, and can function as a cellular transcription factor. The encoded protein can activate transcription through pyrimidine-rich initiator (Inr) elements and E box motifs. USF1 has been linked to familial combined hyperlipidemia (FCHL). The USF1 exists two isoform and the calculated molecular weight of USF1 is a 32 kDa and 27 kDa. USF1 has an acetylated form and phosphorylated form, so the molecular weight of USF1 can migrate to about 40 kDa

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67114-1-Ig (USF1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.