For Research Use Only

ABCA7 Monoclonal antibody

Catalog Number:67128-1-lg



Basic Information

Catalog Number: GenBank Accession Number:

67128-1-lg NM 019112 GeneID (NCBI): Size:

150ul, Concentration: 2500 ug/ml by 10347 2E1A10 Nanodrop and 1000 ug/ml by Bradford_{UNIPROT ID:} Recommended Dilutions: method using BSA as the standard; Q8IZY2

Source: Full Name: Mouse ATP-binding cassette, sub-family A

Isotype: (ABC1), member 7 lgG1 Calculated MW: Immunogen Catalog Number: 234 kDa AG27499 Observed MW:

250 kDa

Applications

Tested Applications: WB, IHC, IF-P, ELISA

Species Specificity: human, mouse, rat, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: 37°C incubated rat brain tissue, Jurkat cells, mouse brain tissue, pig brain tissue, pig thymus tissue, unboiled mouse brain tissue, unboiled rat brain tissue

Purification Method:

WB 1:5000-1:50000

IHC 1:50-1:500

IF-P 1:200-1:800

CloneNo.:

Protein A purification

IHC: human thymus tissue, IF-P: human tonsillitis tissue,

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human thymus tissue slide using 67128-1-Ig (ABCA7 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using ABCA7 antibody (67128-1-Ig, Clone: 2E1A10) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using ABCA7 antibody (67128-1-lg, Clone: 2E1A10) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Various lysates were subjected to SDS PAGE followed by western blot with 67128-1-lg (ABCA7 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.