For Research Use Only

STXBP1 Monoclonal antibody

Catalog Number:67137-1-lg 1 Publications


Antibodies | ELISA kits | Proteins WWW.ptglab.com

Basic Information	Catalog Number: 67137-1-lg	GenBank Accession Number: BC015749		Purification Method: Protein A purification	
	Size:	GenelD (NCBI):		CloneNo.:	
	150ul , Concentration: 2500 ug/ml by			1B5B3	
	Nanodrop and 1000 ug/ml by Bradfor method using BSA as the standard;			Recommended Dilutions: WB 1:5000-1:50000 IHC 1:500-1:2000	
	Source:				
	Mouse	Tutt Name.		IF-P 1:200-1:800	
	lsotype: IgG2a	Calculated MW: 594 aa, 68 kDa			
	Immunogen Catalog Number:Observed MW:AG2860568 kDa				
Applications	Tested Applications:	Tested Applications:Positive Controls:WB, IHC, IF-P, FC (Intra), ELISAWB : fetal human brain tissue, Y79 cells, pig brainCited Applications:tissue, rat brain tissue, mouse brain tissue, pigWBcerebellum tissue, rat cerebellum tissue, mouse			
				brain tissue, mouse brain tissue, pig	
	Species Specificity:	cerebellum		ssue	
	human, mouse, rat, pig	IHC : mouse t		orain tissue, mouse cerebellum tissue	
	Cited Species: human	IF-P : mouse brain tissue,			
	Note-IHC: suggested antigen r TE buffer pH 9.0; (*) Alternativ retrieval may be performed w buffer pH 6.0	vely, antigen			
Background Information	in the regulation of synaptic vesicle of essential for neurotransmission and b	docking and fusion, possil pinds syntaxin, a compon h syntaxins 1, 2, and 3 bu	oly through i ent of the syı	-18/SEC 1 family. STXBP1 may particip nteraction with GTP-binding proteins. I naptic vesicle fusion machinery probat n 4. It may play a role in determining t	
	in the regulation of synaptic vesicle of essential for neurotransmission and b in a 1:1 ratio. STXBP1 can interact wit specificity of intracellular fusion reac	docking and fusion, possil pinds syntaxin, a compon h syntaxins 1, 2, and 3 bu	oly through i ent of the syn t not syntaxi	nteraction with GTP-binding proteins. I naptic vesicle fusion machinery probal	
Background Information Notable Publications	in the regulation of synaptic vesicle of essential for neurotransmission and b in a 1:1 ratio. STXBP1 can interact wit specificity of intracellular fusion read Author Put	docking and fusion, possil binds syntaxin, a compon h syntaxins 1, 2, and 3 bu tions.	oly through i ent of the syn t not syntaxi	nteraction with GTP-binding proteins. I naptic vesicle fusion machinery probat n 4. It may play a role in determining t	
	in the regulation of synaptic vesicle of essential for neurotransmission and b in a 1:1 ratio. STXBP1 can interact wit specificity of intracellular fusion read Author Put	docking and fusion, possil pinds syntaxin, a compon h syntaxins 1, 2, and 3 bu ttions. omed ID Journal 443000 BMC Bio er shipment. % glycerol pH 7.3.	oly through i ent of the syn t not syntaxi	nteraction with GTP-binding proteins. I naptic vesicle fusion machinery probal n 4. It may play a role in determining t Application	


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67137-1-lg (STXBP1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 67137-1-Ig (STXBP1 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 67137-1-Ig (STXBP1 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using STXBP1 antibody (67137-1-Ig, Clone: 1B5B3) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using STXBP1 antibody (67137-1-Ig, Clone: 1B5B3) at dilution of 1:400 and Coralite®488-Conjugated Goat Anti-Mouse IgC(H+L).



1X10^6 HeLa cells were intracellularly stained with 0.4 ug Anti-Human STXBP1 (67137-1-1g, Clone:1B5B3) and CoraLite®488-Conjugated Goat Anti-Mouse 1gG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse 1gG2a 1sotype Control (C1.18.4) (65208-1-1g, Clone: C1.18.4) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).