

For Research Use Only

HDAC3 Monoclonal antibody

Catalog Number: 67151-1-Ig **2 Publications**



Basic Information

Catalog Number: 67151-1-Ig	GenBank Accession Number: BC000614	Purification Method: Protein A purification
Size: 150ul , Concentration: 2000 µg/ml by Nanodrop and 1000 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 8841	CloneNo.: 3E8F10
Source: Mouse	Full Name: histone deacetylase 3	Recommended Dilutions: WB 1:1000-1:6000
Isotype: IgG2b	Calculated MW: 49 kDa	
Immunogen Catalog Number: AG28464	Observed MW: 49 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : HeLa cells, HSC-T6 cells, HEK-293 cells, Jurkat cells
Cited Applications: IP, WB	
Species Specificity: Human, mouse, rat	
Cited Species: human	

Background Information

Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. Histone deacetylase (HDAC) and histone acetyltransferase (HAT) are enzymes that regulate transcription by selectively deacetylating or acetylating the (-amino groups of lysines located near the amino termini of core histone proteins. At least 4 classes of HDAC were identified. HDAC3 is a class I HDAC. HDAC3 has histone deacetylase activity and may participate in the regulation of transcription through its binding with the zinc-finger transcription factor YY1. HDAC3 can also down-regulate p53 function and thus modulate cell growth and apoptosis. The gene encoding HDAC3 is regarded as a potential tumor suppressor gene.

Notable Publications

Author	Pubmed ID	Journal	Application
Xintian Lan	37764423	Molecules	WB
Liu Song	36758802	J Biol Chem	WB,IP

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.1% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67151-1-Ig (HDAC3 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.