For Research Use Only

## ENO1 Monoclonal antibody, PBS Only

Catalog Number:67187-1-PBS

Featured Product





## Catalog Number: GenBank Accession Number: **Purification Method: Basic Information** 67187-1-PBS BC015641 Thiophilic affinity chromatograph GenelD (NCBI): CloneNo.: Size: 100ug , Concentration: 1mg/ml by 2023 1D6E8 Nanodrop: ENSEMBL Gene ID: Source: ENSG0000074800 Mouse UNIPROT ID: Isotype: P06733 IgM Full Name: Immunogen Catalog Number: enolase 1, (alpha) AG17518 Calculated MW: 47 kDa Observed MW: 47 kDa **Applications Tested Applications:** WB, IF/ICC, Indirect ELISA **Species Specificity:** human, mouse, rat, pig **Background Information** ENO1, also named as NNE, ENO1L1, MBPB1, MPB1 and MBP1, belongs to the enolase family. ENO1 is a metabolic enzyme involved in the synthesis of pyruvate. It also acts as a plasminogen receptor and mediates the activation of plasmin and extracellular matrix degradation. In tumor cells, ENO1 is up-regulated and supports the Warburg effect; it is expressed at the cell surface, where it promotes cancer invasion, and is subjected to a specific array of posttranslational modifications, namely acetylation, methylation and phosphorylation. ENO1 overexpression and posttranslational modifications could be of diagnostic and prognostic value in many cancer types. (PMID: 27814656) Storage Storage: Store at -80°C. Storage Buffer: PBS Only

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data





WB result of ENO1 antibody (67187-1-Ig; 1:5000; incubated at room temperature for 1.5 hours) with sh-Control and sh-ENO1 transfected HEK-293 cells. This data was developed using the same antibody clone with 67187-1-PBS in a different storage buffer formulation.

Various lysates were subjected to SDS PAGE followed by western blot with 67187-1-Ig (ENO1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67187-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using ENO1 antibody (67187-1-Ig, Clone: 1D6E8) at dilution of 1:800 and CoraLite® 488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1). This data was developed using the same antibody clone with 67187-1-PBS in a different storage buffer formulation.