For Research Use Only

PITX2 Monoclonal antibody

Catalog Number:67201-1-lg Featured Product

2 Publications



Basic Information

Applications

Catalog Number:

GenBank Accession Number:

Purification Method: Protein A purification

67201-1-lg

GeneID (NCBI):

CloneNo.:

150ul, Concentration: 1700 µg/ml by 5308

BC013998

1E3B1

Nanodrop and 800 µg/ml by Bradford Full Name:

method using BSA as the standard;

paired-like homeodomain 2

Recommended Dilutions: WB 1:5000-1:50000

Calculated MW: 36 kDa

Mouse Isotype: lgG1

Observed MW: 30 kDa

Immunogen Catalog Number:

AG27924

Tested Applications: WB. FIISA

Cited Applications: IF, IHC, WB

Species Specificity:

Human, Mouse, Rat

Cited Species:

human, rat, mouse

Positive Controls:

WB: rat brain tissue, HEK-293 cells, mouse brain tissue, rat cerebellum tissue, mouse cerebellum tissue

Background Information

PITX2 belongs to the bicoid class of homeodomain transcription factors, which play essential roles in embryonic development. Various PITX2 isoforms are involved in a wide variety of developmental programs, including leftright signaling. It controls cell proliferation in a tissue-specific manner and is involved in morphogenesis. During embryonic development, PITX2 is involved in the expansion of muscle progenitors. In addition, it may play a role in the proper localization of asymmetric organs such as the heart and stomach. PITX2 can be bound by some other proteins, and are part of a much larger atrial transcriptional complex. This antibody specifically recognizes isoform PTX2C.

Notable Publications

Author	Pubmed ID	Journal	Application
Yi Yang	35836351	Oral Dis	WB,IHC,IF
Heng Zhang	37950730	Aging (Albany NY)	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67201-1-lg (PITX2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Beta Actin Monoclonal antibody (HRP-66009) as loading control.



HEK-293 cells were subjected to SDS PAGE followed by western blot with 67201-1-1g (PITX2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.