

For Research Use Only

EIF5A Monoclonal antibody

Catalog Number: 67214-1-Ig



Basic Information

Catalog Number: 67214-1-Ig	GenBank Accession Number: BC001832	Purification Method: Protein G purification
Size: 150ul , Concentration: 1000 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 1984	CloneNo.: 2D1B6
Source: Mouse	Full Name: eukaryotic translation initiation factor 5A	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG1	Calculated MW: 18 kDa	
Immunogen Catalog Number: AG28588	Observed MW: 18 kDa	

Applications

Tested Applications:

WB, ELISA

Species Specificity:

Human, mouse, rat

Positive Controls:

WB : HeLa cells, HEK-293T cells, HEK-293 cells, PC-3 cells, Jurkat cells, HSC-T6 cells, NIH/3T3 cells, RAW 264.7 cells

Background Information

Translation of messenger RNA (mRNA) to protein in eukaryotes is a crucial process in protein biosynthesis, in which initiation of translation involves interaction of different eukaryotic translation initiation factors (eIFs), ribosome subunits and mRNAs. Eukaryotic translation initiation factor 5A (EIF5A) is one of the eIFs involved in translation initiation. EIF5A has an essential role in cell viability and is the only protein known to contain the amino acid residue hypusine, formed by post-translational modification of a specific lysine residue. First identified as a translation initiation factor, it also has a function in the elongation step of translation.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67214-1-Ig (EIF5A antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.