## For Research Use Only

## CDK9 Monoclonal antibody

Catalog Number: 67256-1-Ig



**Basic Information** 

Catalog Number: GenBank Accession Number:

67256-1-lg BC001968 GeneID (NCBI): Size:

150ul, Concentration: 1000 ug/ml by 1025 Nanodrop: **UNIPROT ID:** P50750 Mouse Full Name:

Isotype: cyclin-dependent kinase 9

lgG1 Calculated MW: Immunogen Catalog Number: 372 aa, 43 kDa AG29178 Observed MW: 38-42 kDa, 55 kDa **Purification Method:** Protein A purification

CloneNo.:

1G6B3

Recommended Dilutions: WB 1:5000-1:50000 IF/ICC 1:500-1:2000

**Applications** 

**Tested Applications:** 

WB, IF/ICC, ELISA WB: HeLa cells, T-47D cells, HCT 116 cells, A431 cells, Species Specificity: MOLT-4 cells, TF-1 cells, HSC-T6 cells, ROS1728 cells Human, Mouse, Rat

Positive Controls:

**Background Information** 

CDK9(Cyclin-dependent kinase 9) is a member of the Cdc2-like family of kinases. Its cyclin partners are members of the family of cyclin T (T1, T2a and T2b) and cyclin K. The CDK9/cyclin T complexes appear to be involved in regulating several physiological processes. CDK9 has also been described as the kinase of the TAK complex, which is homologous to the P-TEFb complex and involved in HIV replication. In addition, CDK9 seems to have an antiapoptotic function in monocytes, that may be related to its control over differentiation of monocytes (PMID: 12432243). CDK9 has two isoforms with the molecular mass of 42 kDa and 55 kDa, and the relative abundance of Cdk9(42kDa) and Cdk9(55kDa) changes in different cell types (PMID: 12706900, 15780980).

Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.1% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


\*\*\* 20ul sizes contain 0.1% BSA

in USA), or 1(312) 455-8498 (outside USA)

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67256-1-1g (CDK9 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using CDK9 antibody (67256-1-Ig, Clone: 1G6B3) at dilution of 1:1000 and Multi-rAbCoraLite Plus 594-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (RGAM004, 1:500) were used for detection.