## For Research Use Only

## ACSM5 Monoclonal antibody

Catalog Number:67334-1-lg 1 Publications



**Purification Method:** 

WB 1:5000-1:50000

IF/ICC 1:400-1:1600

WB: 4T1 cells, MDA-MB-231 cells, rat liver tissue

IHC 1:250-1:1000

Positive Controls:

IF/ICC: HepG2 cells,

IHC: human liver cancer tissue,

**Basic Information** 

Catalog Number: GenBank Accession Number:

67334-1-lg BC016703 Protein G purification GeneID (NCBI): Size: CloneNo.:

150ul , Concentration: 1800 ug/ml by 549882D5D6 Nanodrop and 1000 ug/ml by Bradford<sub>UNIPROT ID:</sub> Recommended Dilutions: method using BSA as the standard; Q6NUN0

Source: Full Name: Mouse acyl-CoA synthetase medium-chain

Isotype: family member 5 lgG1 Calculated MW: 208 aa, 23 kDa Immunogen Catalog Number: AG10056 Observed MW:

60-65 kDa

**Applications** 

**Tested Applications:** WB, IHC, IF/ICC, ELISA

**Cited Applications:** 

Species Specificity: Human, Mouse, Rat **Cited Species:** 

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

**Background Information** 

ACSM5(Acyl-CoA synthetase medium-chain family member 5) is also named as MACS3 and belongs to the ATPdependent AMP-binding enzyme family. ACSM5 has medium-chain fatty acid:CoA ligase activity with broad substrate specificity (in vitro). This protein has 2 isoforms produced by alternative splicing.

**Notable Publications** 

Author **Pubmed ID** Journal Application Yanlin Cao 34593020 Arthritis Res Ther WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## Selected Validation Data




4T1 cells were subjected to SDS PAGE followed by western blot with 67334-1-1g (ACSM5 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.




rat liver tissue were subjected to SDS PAGE followed by western blot with 67334-1-1g (ACSM5 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 67334-1-Ig (ACSM5 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 67334-1-lg (ACSM5 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using ACSM5 antibody (67334-1-Ig, Clone: 2D5D6) at dilution of 1:800 and CoraLite® 488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1), CL594-phalloidin (red).