## For Research Use Only

## MLH1 Monoclonal antibody

Catalog Number:67350-1-lg Featured Product



**Basic Information** 

Catalog Number: GenBank Accession Number:

67350-1-lg BC006850 Protein A purification GeneID (NCBI): CloneNo.: Size:

150ul, Concentration: 2100 µg/ml by 4292 2F12C4 Nanodrop and 1000 µg/ml by Bradford<sub>Full Name</sub>: Recommended Dilutions:

mutL homolog 1, colon cancer, nonpolyposis type 2 (E. coli)

Mouse Calculated MW: 756 aa, 85 kDa Isotype: lgG2a Observed MW:

method using BSA as the standard;

Immunogen Catalog Number: 85-100 kDa, 40-45 kDa

AG27723

**Applications** 

**Tested Applications:** 

WB, ELISA

Species Specificity: Human, rat

**Positive Controls:** 

WB: A431 cells, HeLa cells, Caco-2 cells, COLO 320

**Purification Method:** 

WB 1:5000-1:50000

cells, HEK-293 cells, Jurkat cells

## **Background Information**

MLH1, also named as COCA2, belongs to the DNA mismatch repair mutL/hexB family. It heterodimerizes with PMS2 to form MutL alpha which is a component of the post-replicative DNA mismatch repair system (MMR). MutL alpha (MLH1-PMS2) interacts physically with the clamp loader subunits of DNA polymerase III, suggesting that it may play a role to recruit the DNA polymerase III to the site of the MMR. MLH1 also implicated in DNA damage signaling, a process which induces cell cycle arrest and can lead to apoptosis in case of major DNA damages. MLH1 heterodimerizes with MLH3 to form MutL gamma which plays a role in meiosis.(PMID: 16873062, PMID: 18206974) Defects in MLH1 are the cause of hereditary non-polyposis colorectal cancer type 2 (HNPCC2). Defects in MLH1 are a cause of mismatch repair cancer syndrome (MMRCS). Defects in MLH1 are a cause of Muir-Torre syndrome (MTS). Defects in MLH1 are a cause of susceptibility to endometrial cancer. Western blot analysis with an MLH1 antibody detected a 85-100 kDa band. Full-length human MLH1 is specifically cleaved into degradation products of 40-45 kDa by caspase-3 (PMID: 15087450, PMID: 19603033). This antibody is specific to MLH1.

Storage


Storage: Store at -20°C. Storage Buffer:


PBS with 0.1% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 67350-1-1g (MLH1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.

WB result of MLH1 antibody (67350-1-lg; 1:10000; incubated at room temperature for 1.5 hours) with sh-Control and sh-MLH1 transfected HeLa cells.