For Research Use Only

HMG20B Monoclonal antibody

Catalog Number:67354-1-lg




Basic Information	Catalog Number: 67354-1-1g	GenBank Accession Num BC002552	ber: Purification Method: Protein G purification
	Size: 150ul , Concentration: 1600 ug/ml by		CloneNo.: 4C2A9
	Nanodrop and 1000 ug/ml by Bradfor method using BSA as the standard;	d <mark>UNIPROT ID:</mark> Q9POW2	Recommended Dilutions: WB 1:5000-1:50000
	Source: Mouse	Full Name: high-mobility group 20B	IHC 1:1000-1:4000
	lsotype: IgG1	Calculated MW: 36 kDa	
	Immunogen Catalog Number: AG6103	Observed MW: 40 kDa	
Applications	Tested Applications:	Po	ositive Controls:
	Species Specificity: cells, K-56		B : LNCaP cells, HeLa cells, HEK-293 cells, Jurkat ells, K-562 cells, HSC-T6 cells, NIH/3T3 cells
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0		
Background Information	CoREST (corepressor of repressor element-1-silencing transcription factor) is a chromatin- modifying complex that was first identified as a regulator of neuronal gene expression, having a essential role in neuronal differentiation [PMID:10449787]. The CoREST complex is recruited by REST (RE1-silencing transcription factor) that recognizes the 21-nucleotide RE-1 sequence in the promoter of more than 1,700 different target genes in human and mouse [PMID:15240883]. HMGB20B is one component of the CoREST complex, which mediates chromatin remodeling and repression of specific genes involved in neuronal differentiation through modulation of chromatin structure [PMID:19997604].		
Storage	Storage: Store at -20°C. Stable for one year aft Storage Buffer: PBS with 0.02% sodium azide and 50	% glycerol pH 7.3.	
*** 20ul sizes contain 0.1% BSA	Aliquoting is unnecessary for -20 $^{\circ}$ C s	torage	

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 67354-1-lg (HMG20B antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded human skin cancer tissue slide using 67354-1-Ig (HMG20B antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).