For Research Use Only

TBXAS1 Monoclonal antibody

Catalog Number:67371-1-lg

Featured Product

1 Publications



Basic Information

Catalog Number: GenBank Accession Number: BC014117

67371-1-lg Protein A purification GeneID (NCBI): Size: CloneNo.:

150ul, Concentration: 1900 ug/ml by 6916 1F6G6 Nanodrop and 1000 ug/ml by Bradford_{UNIPROT ID:} Recommended Dilutions: method using BSA as the standard; P24557 WB 1:5000-1:50000

Full Name: Mouse thromboxane A synthase 1 (platelet)

Isotype: Calculated MW: IgG2a 60 kDa Immunogen Catalog Number: Observed MW: AG29177 52 kDa

Applications

Tested Applications:

WB, IHC, FC (Intra), ELISA

Cited Applications:

WB, IHC

Source:

Species Specificity: human, mouse, pig Cited Species:

mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: human peripheral blood platelets, A549 cells,

Purification Method:

IHC 1:1000-1:4000

human placenta tissue, pig blood IHC: human lung cancer tissue,

Background Information

TBXAS1 catalyzes the conversion of the prostaglandin endoperoxide into thromboxane A2 as a potent vasoconstrictor and inducer of platelet aggregation. It was identified as novel airway fibroblast-specific markers with integrin-8 (ITGA8) (PMID:20435685). It was observed in the epithelium. It is a monomer with the molecular weight of 60 KDa. It also has two variants with 60.5 kDa and 52.3 kDa (PMID:1730669). It also reavels a molecular weight of 55 kDa in human uterus (PMID:1386258).

Notable Publications

Author	Pubmed ID	Journal	Application
Katharina Mederle	26017977	Am J Physiol Renal Physiol	WB,IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.1% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Human peripheral blood platelets were subjected to SDS PAGE followed by western blot with 67371-1-Ig (TBXAS1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.




WB result of TBXAS1 antibody (67371-1-1g; 1:5000; incubated at room temperature for 1.5 hours) with sh-Control and sh-TBXAS1 transfected A549 cells.



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 67371-1-1g (TBXAS1 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 67371-1-Ig (TBXAS1 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1x10^6 A549 cells were intracellularly stained with 0.8 ug TBXA51 Monoclonal antibody (67371-1-1g, Clone:1F6G6) and CoraLite® 488-Conjugated Goat Anti-Mouse 1gG(H+L) (SA00013-1)(red), or 0.8 ug Mouse 1gG2a isotype control Mouse McAb (66360-2-1g, Clone: 11A1B2) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).