

For Research Use Only

UBAC1 Monoclonal antibody

Catalog Number: 67385-1-Ig **1 Publications**



Basic Information

Catalog Number: 67385-1-Ig	GenBank Accession Number: BC004967	Purification Method: Protein G purification
Size: 150ul , Concentration: 2100 µg/ml by Nanodrop and 1000 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 10422	CloneNo.: 2A3G7
Source: Mouse	UNIPROT ID: Q9BSL1	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG1	Full Name: UBA domain containing 1	
Immunogen Catalog Number: AG13958	Calculated MW: 405 aa, 45 kDa	
	Observed MW: 46 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : LNCaP cells, HeLa cells, Jurkat cells, K-562 cells, pig brain tissue, mouse brain tissue, rabbit brain tissue, rat brain tissue
Cited Applications: WB, ColP	
Species Specificity: Human, Mouse, Rat, pig, rabbit	
Cited Species: human	

Background Information

UBAC1(Ubiquitin-associated domain-containing protein 1) is also named as GBDR1, KPC2, UBADC1. It has several putative phosphorylation sites, including 8 CK2 (see 115440) sites, as well as 1 N-glycosylation site and 2 N-myristoylation sites(PMID:10857748). It is required for poly-ubiquitination and proteasome-mediated degradation of CDKN1B during G1 phase of the cell cycle.Northern blot analysis shows UBAC1 expression with the highest levels in heart and skeletal muscle and lowest level in lung.

Notable Publications

Author	Pubmed ID	Journal	Application
Hanqi Li	38479395	Cell Host Microbe	WB,ColP

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67385-1-Ig (UBAC1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.