

GRIM19 Monoclonal antibody

Catalog Number: 67414-1-Ig

Basic Information

Catalog Number: 67414-1-Ig	GenBank Accession Number: BC009189	Purification Method: Protein G purification
Size: 150ul , Concentration: 1100 ug/ml by Nanodrop and 1000 ug/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 51079	CloneNo.: 3A10C5
Source: Mouse	UNIPROT ID: Q9P0J0	Recommended Dilutions: WB 1:1000-1:6000 IHC 1:500-1:2000
Isotype: IgG1	Full Name: NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 13	
Immunogen Catalog Number: AG20855	Calculated MW: 16.5 kDa	
	Observed MW: 16-18 kDa	

Applications

Tested Applications: WB, IHC, ELISA	Positive Controls:
Species Specificity: Human	WB : LNCaP cells, HEK-293 cells, HeLa cells, HepG2 cells
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	IHC : human liver cancer tissue, human breast cancer tissue

Background Information

GRIM19(Gene associated with retinoic and interferon-induced mortality 19 protein) is also named as NDUFA13, CI-B16.6 and belongs to the complex I NDUFA13 subunit family. GRIM-19 may distribute in the mitochondria and/or nucleus depending on the cell type. Alternatively, its cellular localization may be regulated by post-translational modifications such as phosphorylation and acetylation. Lastly, GRIM-19 may exist in different isoforms that may be distributed in distinct cellular locations(PMID:12628925).

Storage


Storage:
Store at -20°C.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

***** 20ul sizes contain 0.1% BSA**


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




LNCaP cells were subjected to SDS PAGE followed by western blot with 67414-1-Ig (GRIM19 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



HepG2 cells were subjected to SDS PAGE followed by western blot with 67414-1-Ig (GRIM19 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 67414-1-Ig (GRIM19 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 67414-1-Ig (GRIM19 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).