

For Research Use Only

# MAP1B Monoclonal antibody

Catalog Number: 67423-1-Ig



## Basic Information

<b>Catalog Number:</b> 67423-1-Ig	<b>GenBank Accession Number:</b> BC141853	<b>Purification Method:</b> Protein G purification
<b>Size:</b> 150ul , Concentration: 2100 µg/ml by Nanodrop and 1000 µg/ml by Bradford method using BSA as the standard;	<b>GeneID (NCBI):</b> 4131	<b>CloneNo.:</b> 1E6B9
<b>Source:</b> Mouse	<b>Full Name:</b> microtubule-associated protein 1B	<b>Recommended Dilutions:</b> IHC 1:2500-1:10000 IF 1:50-1:500
<b>Isotype:</b> IgG1	<b>Calculated MW:</b> 2468 aa, 271 kDa	
<b>Immunogen Catalog Number:</b> AG17709		

## Applications

<b>Tested Applications:</b> FC, IF, IHC, ELISA	<b>Positive Controls:</b> IHC : rat brain tissue, mouse brain tissue IF : rat brain tissue,
<b>Species Specificity:</b> Human, rat, mouse	
<b>Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0</b>	

## Background Information

Microtubule-associated protein 1B (MAP1B) is a cytoskeleton protein which can promote microtubule assembly. Previous reports have suggested that this protein is closely involved in neuronal development based on its extensive expression in the developing brain and moderate in mature neurons. Gene disruption or knockout studies of the MAP1B gene led to a delayed development of the nervous system in mice. It includes the N-terminal heavy chain and a C-terminal light chain. The MAP1B heavy chain has a microtubule-stabilization effect, and contains an actin-binding site that may play a role in the crosslinking of actin and microtubules, a function that may be important in neurite elongation. Various isoforms around 300-350 kDa of MAP1B can be observed due to the differences in phosphorylation state. (10704485)

## Storage


**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.1% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Immunohistochemical analysis of paraffin-embedded rat brain tissue slide using 67423-1-Ig (MAP1B antibody) at dilution of 1:5000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed rat brain tissue using 67423-1-Ig (MAP1B antibody), at dilution of 1:200 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunohistochemical analysis of paraffin-embedded rat brain tissue slide using 67423-1-Ig (MAP1B antibody) at dilution of 1:5000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1X10<sup>6</sup> SH-SY5Y cells were intracellularly stained with 0.4 ug Anti-Human MAP1B (67423-1-Ig, Clone:1E6B9) and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG1 Isotype Control (MOPC-21) (65124-1-Ig, Clone: MOPC-21) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).