For Research Use Only

PROX1 Monoclonal antibody

Catalog Number:67438-1-lg 1 Publications



Basic Information

Catalog Number: GenBank Accession Number:

67438-1-lg BC024201 GeneID (NCBI): Size:

150ul , Concentration: 2000 $\mu g/ml$ by 5629 Nanodrop and 1000 µg/ml by Bradford_{UNIPROT ID:} method using BSA as the standard; Q92786

Source: Full Name: Mouse

prospero homeobox 1 Isotype: Calculated MW: lgG2b 737 aa, 83 kDa Immunogen Catalog Number: Observed MW: AG9681 90 kDa

Purification Method:

Protein A purification

CloneNo.: 1D4B1

Recommended Dilutions:

WB 1:1000-1:6000 IHC 1:2000-1:8000 IF 1:200-1:800

Applications

Tested Applications: IF, IHC, WB, ELISA

Species Specificity: Human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: HuH-7 cells, rat liver tissue IHC: human lung cancer tissue,

IF: HuH-7 cells,

Background Information

PROX1, Prospero homeobox protein 1, belongs to the Prospero homeobox family and contain a Prospero-type homeobox DNA-binding domain. PROX1 plays a fundamental role in early development of CNS and regulates the expression of gene and development of postmitotic, undifferentiated neurons. Also, PROX1 is a required regulator of the development of the lymphatic system. This antibody is raised against part of C-terminal chain of human PROX1. The calculated molecular weight of PROX1 is 83 kDa, but with post-translational modifications, PROX1 is about 90

Notable Publications

Author	Pubmed ID	Journal	Application
Wenxiu Dai	38272027	Dev Cell	

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

*** 20ul sizes contain 0.1% BSA

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




HuH-7 cells were subjected to SDS PAGE followed by western blot with 67438-1-1g (PROX1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 67438-1-1g (PROX1 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 67438-1-Ig (PROX1 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HuH-7 cells using PROX1 antibody (67438-1-lg, Clone: 1D4B1) at dilution of 1:400 and CoraLite®488-Conjugated Affini Pure Goat Anti-Mouse IgG(H+L).