

For Research Use Only

CCT4 Monoclonal antibody

Catalog Number: 67455-1-Ig



Basic Information

Catalog Number:

67455-1-Ig

Size:

150ul, Concentration: 1000 ug/ml by Nanodrop and 625 ug/ml by Bradford method using BSA as the standard;

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG17084

GenBank Accession Number:

BC106934

GeneID (NCBI):

10575

UNIPROT ID:

P50991

Full Name:

chaperonin containing TCP1, subunit 4 (delta)

Calculated MW:

539 aa, 58 kDa

Observed MW:

55-58 kDa

Purification Method:

Protein G purification

CloneNo.:

3C6A6

Recommended Dilutions:

WB 1:5000-1:50000

IF/ICC 1:200-1:800

Applications

Tested Applications:

WB, IF/ICC, ELISA

Species Specificity:

Human, Mouse, Rabbit, Pig

Positive Controls:

WB : LNCaP cells, HeLa cells, HEK-293 cells, HepG2 cells, Jurkat cells, pig brain tissue, rabbit brain tissue, mouse brain tissue

IF/ICC : U2OS cells,

Background Information

T-complex protein 1 subunit delta (CCT4) is a component of the chaperonin-containing T-complex (TRiC), a molecular chaperone complex that assists the folding of proteins upon ATP hydrolysis (PMID:25467444). As part of the TRiC complex, it may play a role in regulating telomere maintenance (PMID:25467444) and assembly of BBSome, a complex involved in ciliogenesis regulating transports vesicles to the cilia (PMID:20080638). CCT4 has 2 isoforms produced by alternative splicing with the MW of 55 kDa and 58 kDa.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.1% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67455-1-Ig (CCT4 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.



Immunofluorescent analysis of (-20°C Methanol) fixed U2OS cells using CCT4 antibody (67455-1-Ig, Clone: 3C6A6) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L).