For Research Use Only

CMPK1 Monoclonal antibody

Catalog Number: 67519-1-Ig



Purification Method:

Basic Information

Catalog Number: GenBank Accession Number:

67519-1-lg BC014961 Protein A purification

Size: GeneID (NCBI): CloneNo.: 150ul , Concentration: 1000 µg/ml by 51727 2D11G7

Nanodrop; Full Name: Recommended Dilutions:
Source: cytidine monophosphate (UMP-CMP) WB 1:5000-1:50000
Mouse kinase 1, cytosolic IF 1:200-1:800

Isotype:Calculated MW:IgG2b228 aa, 26 kDaImmunogen Catalog Number:Observed MW:AG3008422-26 kDa

Applications

Tested Applications: Positive Controls:

IF, WB, ELISA WB : HeLa cells, HEK-293 cells, Jurkat cells

Species Specificity: IF : HeLa cells,

Background Information

Pyrimidine nucleoside monophosphate kinase [UMP/CMP kinase (UMP/CMPK)] (CMPK1) is also named as CMK, CMPK, UCK, UMK, UMPK and belongs to the adenylate kinase family. It plays a crucial role in the formation of UDP, CDP, and dCDP, which are required for cellular nucleic acid synthesis(PMID:11912132). CMPK1 can use a broad spectrum of phosphate donors but the best phosphate donors are ATP and dATP. CMPK1 has been reported as a prognostic marker for non-small cell lung cancer, pancreatic cancer and breast cancer (PMID: 21642870, 22838950, 27558661). The longest deduced protein contains 228 amino acids and has a calculated molecular mass of 26 kD. However, translation likely begins at the second initiation methionine, resulting in a protein of 196 amino acids(PMID:10462544). The shorter form is the endogenously translated protein(PMID:11912132). CMPK1 has 3 isoforms with the molecular mass of 22, 16 and 26 kDa.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67519-1-1g (CMPK1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using CMPK1 antibody (67519-1-lg, Clone: 2D11G7) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).