

For Research Use Only

UAP1 Monoclonal antibody

Catalog Number: 67545-1-Ig

Featured Product

3 Publications



Basic Information

Catalog Number:

67545-1-Ig

Size:

150ul, Concentration: 2400 ug/ml by 6675
Nanodrop and 1000 ug/ml by Bradford
method using BSA as the standard;

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG9513

GenBank Accession Number:

BC009377

GeneID (NCBI):

6675

UNIPROT ID:

Q16222

Full Name:

UDP-N-acetylglucosamine
pyrophosphorylase 1

Calculated MW:

505 aa, 57 kDa

Observed MW:

57-59 kDa

Purification Method:

Protein A purification

CloneNo.:

2C7B2

Recommended Dilutions:

WB 1:5000-1:50000

IHC 1:500-1:2000

IF/ICC 1:200-1:800

Applications

Tested Applications:

WB, IHC, IF/ICC, ELISA

Cited Applications:

WB, IHC, CoIP

Species Specificity:

Human

Cited Species:

human

**Note-IHC: suggested antigen retrieval with
TE buffer pH 9.0; (*) Alternatively, antigen
retrieval may be performed with citrate
buffer pH 6.0**

Positive Controls:

WB : HeLa cells, A549 cells, NCI-H1299 cells, LNCaP
cells, HepG2 cells, THP-1 cells, HEK-293 cells, DU 145
cells, Jurkat cells, K-562 cells

IHC : human pancreas cancer tissue,

IF/ICC : PC-3 cells,

Background Information

UAP1 can convert UTP and GlcNAc-1-P into UDP-GlcNAc, it can also convert UTP and GalNAc-1-P into UDP-GalNAc. There are three isoforms of UAP1, among which isoform AGX1 has 2 to 3 times higher activity towards GalNAc-1-P, while isoform AGX2 has 8 times more activity towards GlcNAc-1-P. It has been reported that UAP1 is specific and highly expressed in prostate cancer. UAP1 can protect cells against inhibitors of N-linked glycosylation. UAP1 may play an important role in maintaining INS responsiveness. 67545-1-Ig detects 57 and 59 kDa isoforms in SDS-PAGE. (PMID:25241896)

Notable Publications

Author	Pubmed ID	Journal	Application
Zexin Su	36275679	Front Immunol	IHC, WB
Shuai Yang	36603579	Mol Cell	WB
Yuan-Li Ai	37460805	Cell Res	WB, CoIP

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech
Group brand and is not available to purchase from any
other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67545-1-Ig (UAP1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human pancreas cancer tissue slide using 67545-1-Ig (UAP1 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed PC-3 cells using UAP1 antibody (67545-1-Ig, Clone: 2C7B2) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).