For Research Use Only

NXT1 Monoclonal antibody

Catalog Number: 67680-1-lg 1 Publications



Basic Information

Catalog Number: GenBank Accession Number:

67680-1-lg BC000759

Size: Genel D (NCBI):
150ul , Concentration: 1600 ug/ml by 29107

Nanodrop and 1000 ug/ml by Bradford UNIPROT ID: method using BSA as the standard; O9UKK6

Source: Full Name:
Mouse NTE2-like export

Mouse NTF2-like export factor 1
Isotype: Calculated MW:

IgG2a 15 kDa

Immunogen Catalog Number: Observed MW:
AG30039 15 kDa

Purification Method:

Protein A purification CloneNo.:

2C10A2
Recommended Dilutions:

Recommended Dilutior WB 1:5000-1:50000 IF/ICC 1:200-1:800

Applications

Tested Applications: WB, IF/ICC, ELISA

Cited Applications: WB

Species Specificity: Human, mouse, rat Cited Species: human Positive Controls:

WB: A549 cells, HEK-293 cells, HeLa cells, HepG2 cells,

HSC-T6 cells, NIH/3T3 cells, 4T1 cells

IF/ICC: HeLa cells,

Background Information

NXT1 is located in the nuclear envelope and is homologous to nuclear transport factor 2. NXT1 functions as a nuclear export factor in both RAN (Ras-related nuclear protein)- and CRM1 (required for chromosome region maintenance)-dependent pathways. It is found to stimulate the export of U1 snRNA in RAN- and CRM1-dependent pathways and the export of tRNA and mRNA in a CRM1-independent pathway. NXT1 also heterodimerizes with Tap protein and regulates the ability of Tap protein to mediate nuclear mRNA export.

Notable Publications

Author	Pubmed ID	Journal	Application
Mio Iwasaki	35573189	iScience	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67680-1-lg (NXT1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using NXT1 antibody (67680-1-1g, Clone: 2C10A2) at dilution of 1:400 and Coralite® 488-Conjugated Affini Pure Goat Anti-Mouse IgG(H+L).



Various lysates were subjected to SDS PAGE followed by western blot with 67680-1-1g (NXT1 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.