For Research Use Only

FABP2 Monoclonal antibody

Catalog Number:67691-1-lg Featured Product

2 Publications



Basic Information

Catalog Number: GenBank Accession Number: **Purification Method:** 67691-1-lg BC069617 Protein G purification

GeneID (NCBI): Size: CloneNo.: 150ul, Concentration: 1000 ug/ml by 2169 2D11G6

Nanodrop: **UNIPROT ID:** Recommended Dilutions: P12104 WB 1:2000-1:50000 IHC 1:2000-1:8000 Mouse Full Name: fatty acid binding protein 2, intestinal IF-P 1:200-1:800 Isotype:

lgG1 Calculated MW: Immunogen Catalog Number: 132 aa, 15 kDa AG17620 Observed MW: 15 kDa

Applications

Tested Applications: WB, IHC, IF-P, ELISA

Cited Applications: WB. IF

Species Specificity:

Human, mouse, rat, rabbit, pig

Cited Species:

mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: rat small intestine tissue, human jejunum tissue, COLO 320 cells, pig duodenum, mouse small intestine, rabbit small intestine

IHC: mouse small intestine tissue, mouse colon tissue, rat small intestine tissue, human small intestine

tissue

IF-P: mouse colon tissue.

Background Information

FABP2, also known as the intestinal fatty acid binding protein (I-FABP), is expressed in the absorptive intestinal villus cells. It is mainly involved in intracellular transport and intestinal absorption of lipids. FABP2 has been considered a marker of mucosal injury and ischemia and serum I-FABP level is used as a tissue damage indicator. In addition, it is a marker of differentiated intestinal epithelial cells.

Notable Publications

Author	Pubmed ID	Journal	Application
Li-Long Pan	39648298	Gut Microbes	IF
Yunzhe Su	38003599	Int J Mol Sci	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67691-1-1g (FABP2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (4% PFA) fixed mouse colon tissue using FABP2 antibody (67691-1-Ig, Clone: 2D11G6) at dilution of 1:400 and CoraLite@488-Conjugated Goat Anti-Mouse IgG(H+L).




WB result of FABP2 antibody (67691-1-lg; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-FABP2 transfected COLO 320 cells




Immunohistochemical analysis of paraffinembedded mouse small intestine tissue slide using 67691-1-Ig (FABP2 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse small intestine tissue slide using 67691-1-lg (FABP2 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded rat small intestine tissue slide using 67691-1-lg (FABP2 antibody) at dilution of 1:8000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human small intestine tissue slide using 67691-1-Ig (FABP2 antibody) at dilution of 1:8000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human small intestine tissue slide using 67691-1-1g (FABP2 antibody) at dilution of 1:8000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).